

t73_sprect_1

(TMR8yZn43nfNQkB2EjAMxPYLJatBvteAyxn)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v2_compts_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k7_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $k4_pscomp_1 : \iota$ be given. Let $k1_rltopsp1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k13_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k10_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k1_rcomp_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k8_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k19_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rlvect_1 : \iota \Rightarrow o$ be given. Let $v6_rlvect_1 : \iota \Rightarrow o$ be given. Let $v7_rlvect_1 : \iota \Rightarrow o$ be given. Let $v8_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rltopsp1 : \iota \Rightarrow o$ be given. Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_rltopsp1 : \iota \Rightarrow o$ be given. Let $l1_rlvect_1 : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $k9_pscomp_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\
 & (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\
 & ((r1_xxreal_0 (k17_euclid X0) (k17_euclid X1)) \Rightarrow (k7_relset_1 \\
 & (u1_struct_0 (k15_euclid np_2)) k1_numbers k4_pscomp_1 (k1_rltopsp1 \\
 & (k15_euclid np_2) X0 X1) = k1_rcomp_1 (k17_euclid X0) (k17_euclid \\
 & X1))))
 \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((k17_euclid \\
 (k19_euclid X0 X1) = X0) \wedge (k18_euclid (k19_euclid X0 X1) = X1))) \tag{2}$$

Assume the following.

$$\forall X0.((\neg v1_xboole_0 X0) \wedge ((v2_compts_1 X0 (k15_euclid np_2)) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))))) \Rightarrow (r1_xxreal_0 (k6_pscomp_1 X0) (k8_pscomp_1 X0))) \quad (3)$$

Assume the following.

$$((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \quad (4)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (5)$$

Assume the following.

$$v6_membered k4_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow ((v2_pre_topc (k15_euclid X0)) \wedge ((v13_algstr_0 (k15_euclid X0)) \wedge ((v2_rlvect_1 (k15_euclid X0)) \wedge ((v3_rlvect_1 (k15_euclid X0)) \wedge ((v4_rlvect_1 (k15_euclid X0)) \wedge ((v5_rlvect_1 (k15_euclid X0)) \wedge ((v6_rlvect_1 (k15_euclid X0)) \wedge ((v7_rlvect_1 (k15_euclid X0)) \wedge ((v8_rlvect_1 (k15_euclid X0)) \wedge (v5_rltopsp1 (k15_euclid X0)))))))))))))) \quad (7)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow ((\neg v2_struct_0 (k15_euclid X0)) \wedge (v5_rltopsp1 (k15_euclid X0))) \quad (8)$$

Assume the following.

$$\forall X0.(l1_rltopsp1 X0) \Rightarrow ((l1_rlvect_1 X0) \wedge (l1_pre_topc X0)) \quad (9)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (m1_subset_1 (k9_pscomp_1 X0) k1_numbers) \quad (10)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (m1_subset_1 (k8_pscomp_1 X0) k1_numbers) \quad (11)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (m1_subset_1 (k6_pscomp_1 X0) k1_numbers) \quad (12)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow ((v5_rltopsp1 (k15_euclid X0)) \wedge (l1_rltopsp1 (k15_euclid X0))) \quad (13)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (m1_subset_1 (k13_pscomp_1 X0) (u1_struct_0 (k15_euclid np_2))) \quad (14)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (m1_subset_1 (k10_pscomp_1 X0) (u1_struct_0 (k15_euclid np_2))) \quad (15)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (k13_pscomp_1 X0 = k19_euclid (k8_pscomp_1 X0) (k9_pscomp_1 X0)) \quad (16)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (k10_pscomp_1 X0 = k19_euclid (k6_pscomp_1 X0) (k9_pscomp_1 X0)) \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v2_rlvect_1 X0) \wedge ((v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v5_rlvect_1 X0) \wedge ((v6_rlvect_1 X0) \wedge ((v7_rlvect_1 X0) \wedge ((v8_rlvect_1 X0) \wedge (l1_rlvect_1 X0)))))))))) \wedge ((m1_subset_1 X1 (u1_struct_0 X0)) \wedge (m1_subset_1 X2 (u1_struct_0 X0)))) \Rightarrow (k1_rltopsp1 X0 X1 X2 = k1_rltopsp1 X0 X2 X1) \quad (18)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (19)$$

Assume the following.

$$\forall X0.(v6_membered X0) \Rightarrow (\forall X1.(m1_subset_1 X1 X0) \Rightarrow (v7_ordinal1 X1)) \quad (20)$$

Theorem 1

$$\begin{aligned} & \forall X0. ((\neg v1_xboole_0 X0) \wedge ((v2_compts_1 X0 (k15_euclid np_2)) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))))) \Rightarrow \\ & (k7_relset_1 (u1_struct_0 (k15_euclid np_2)) k1_numbers k4_pscomp_1 \\ & (k1_rltopsp1 (k15_euclid np_2) (k13_pscomp_1 X0) (k10_pscomp_1 \\ & X0)) = k1_rcomp_1 (k6_pscomp_1 X0) (k8_pscomp_1 X0)) \end{aligned}$$