

# t73\_xboole\_1 (TMYQPWYEGdobn- rmL1FqDkAC8L4MpwWwENe1)

October 27, 2020

Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. (r1\_tarski X0 X1) \Rightarrow (X1 = k2\_xboole\_0 X0 (k4\_xboole\_0 X1 X0)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (\neg(\neg r1\_xboole\_0 X0 X1) \wedge (\forall X2. \neg(X2 \in X0) \wedge (X2 \in X1))) \wedge (\neg(\exists X2. (X2 \in X0) \wedge (X2 \in X1)) \wedge (r1\_xboole\_0 X0 X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k2\_xboole\_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \vee (X3 \in X1))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (r1\_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (4)$$

## Theorem 1

$$\forall X0. \forall X1. \forall X2. ((r1\_tarski X0 (k2\_xboole\_0 X1 X2)) \wedge (r1\_xboole\_0 X0 X2)) \Rightarrow (r1\_tarski X0 X1)$$