

# t73\_xboolean (TM- RmYgi4KBqeCGWEASXm3Q7bRtf1BPPuKqG)

October 27, 2020

Let  $v1\_xboolean : \iota \Rightarrow o$  be given. Let  $k10\_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k5\_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_0 : \iota$  be given. Let  $k3\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k6\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k3\_xboolean : \iota \Rightarrow \iota$  be given. Let  $k2\_xboolean : \iota$  be given. Let  $k1\_xboolean : \iota$  be given. Let  $k7\_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(v1\_xboolean X0) \Rightarrow (\forall X1.(v1\_xboolean X1) \Rightarrow (k10\_xboolean X0 (k10\_xboolean X0 X1) = X1)) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (2)$$

Assume the following.

$$\forall X0.(v1\_xboolean X0) \Rightarrow (\forall X1.(v1\_xboolean X1) \Rightarrow (k5\_xboolean X0 (k4\_xboolean X0 X1) = X0)) \quad (3)$$

Assume the following.

$$v1\_xboole\_0 np\_0 \quad (4)$$

Assume the following.

$$k3\_xcmplx\_0 np\_1 np\_0 = np\_0 \quad (5)$$

Assume the following.

$$k6\_xcmplx\_0 np\_1 np\_0 = np\_1 \quad (6)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (7)$$

Assume the following.

$$\exists X0.v1\_xboolean X0 \quad (8)$$

Assume the following.

$$\forall X0.(v1\_xboolean\ X0)\Rightarrow(k3\_xboolean\ (k3\_xboolean\ X0) = X0) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xboolean\ X0)\wedge(v1\_xboolean\ X1))\Rightarrow(k4\_xboolean\ X0\ X0 = X0) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xboolean\ X0)\wedge(v1\_xboolean\ X1))\Rightarrow(v1\_xboolean\ (k10\_xboolean\ X0\ X1)) \quad (11)$$

Assume the following.

$$v1\_xboolean\ k2\_xboolean \quad (12)$$

Assume the following.

$$v1\_xboolean\ k1\_xboolean \quad (13)$$

Assume the following.

$$\forall X0.(v1\_xboolean\ X0)\Rightarrow(\forall X1.(v1\_xboolean\ X1)\Rightarrow(k7\_xboolean\ X0\ X1 = k4\_xboolean\ (k6\_xboolean\ X0\ X1)\ (k6\_xboolean\ X1\ X0))) \quad (14)$$

Assume the following.

$$\forall X0.(v1\_xboolean\ X0)\Rightarrow(\forall X1.(v1\_xboolean\ X1)\Rightarrow(k6\_xboolean\ X0\ X1 = k5\_xboolean\ (k3\_xboolean\ X0)\ X1)) \quad (15)$$

Assume the following.

$$\forall X0.(v1\_xboolean\ X0)\Rightarrow(\forall X1.(v1\_xboolean\ X1)\Rightarrow(k4\_xboolean\ X0\ X1 = k3\_xcmplx\_0\ X0\ X1)) \quad (16)$$

Assume the following.

$$\forall X0.(v1\_xboolean\ X0)\Rightarrow(k3\_xboolean\ X0 = k6\_xcmplx\_0\ np\_1\ X0) \quad (17)$$

Assume the following.

$$\forall X0.(v1\_xboolean\ X0)\Leftrightarrow((X0 = k1\_xboolean)\vee(X0 = k2\_xboolean)) \quad (18)$$

Assume the following.

$$k2\_xboolean = np\_1 \quad (19)$$

Assume the following.

$$k1\_xboolean = k6\_numbers \quad (20)$$

Assume the following.

$$\forall X0.(v1\_xboolean\ X0)\Rightarrow(\forall X1.(v1\_xboolean\ X1)\Rightarrow(k10\_xboolean\ X0\ X1 = k3\_xboolean\ (k7\_xboolean\ X0\ X1))) \quad (21)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xboolean\ X0)\wedge(v1\_xboolean\ X1))\Rightarrow( \\ k5\_xboolean\ X0\ X1 = k5\_xboolean\ X1\ X0) \quad (22)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xboolean\ X0)\wedge(v1\_xboolean\ X1))\Rightarrow( \\ k10\_xboolean\ X0\ X1 = k10\_xboolean\ X1\ X0) \quad (23)$$

**Theorem 1**

$$\forall X0.(v1\_xboolean\ X0)\Rightarrow(\forall X1.(v1\_xboolean\ X1)\Rightarrow(\forall X2. \\ (v1\_xboolean\ X2)\Rightarrow(k10\_xboolean\ (k10\_xboolean\ X0\ X1)\ X2 = k10\_xboolean \\ X0\ (k10\_xboolean\ X1\ X2))))$$