

t74_funcop_1
(TMFFpASPmiuzrquNyD3q7xyPGkf474o5FC6)

October 27, 2020

Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k16_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 \\ X2)) \Rightarrow ((r1_tarski (k16_funcop_1 X0 X1) X2) \Leftrightarrow ((X0 \in k9_xtuple_0 X2) \wedge \\ (k1_funct_1 X2 X0 = X1))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. r1_tarski X0 X0 \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 (k16_funcop_1 X0 X1)) \wedge (v1_funct_1 \\ (k16_funcop_1 X0 X1)) \tag{3}$$

Theorem 1 $\forall X0. \forall X1. X0 \in k9_xtuple_0 (k16_funcop_1 X0 X1).$