

t74_funct_4

(TMZwyYEogtVJXT9sSEkQHQdG9QALquTrSWJ)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((\\ v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_tarski (k9_xtuple_0 X0) \\ (k9_xtuple_0 X1)) \Rightarrow (k1_funct_4 X0 X1 = X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((\\ v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2.((v1_relat_1 X2) \wedge \\ (v1_funct_1 X2)) \Rightarrow (k1_funct_4 (k1_funct_4 X0 X1) X2 = k1_funct_4 \\ X0 (k1_funct_4 X1 X2)))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. r1_tarski X0 X0 \quad (3)$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((\\ v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2.((v1_relat_1 X2) \wedge \\ (v1_funct_1 X2)) \Rightarrow ((k9_xtuple_0 X1 = k9_xtuple_0 X2) \Rightarrow (k1_funct_4 \\ (k1_funct_4 X0 X1) X2 = k1_funct_4 X0 X2)))) \end{aligned}$$