

t74\_lukasi\_1 (TMZy-  
atzE3ZEnRKCHzcGTGG348krhPZrAkTf)

October 27, 2020

Let  $m1\_qc\_lang1 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_qc\_lang1 : \iota \Rightarrow \iota$  be given. Let  $k3\_cqc\_lang : \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $r3\_cqc\_the1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k8\_cqc\_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_cqc\_lang : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m2\_subset\_1 X1 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X2.(m2\_subset\_1 X2 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X3.(m1\_subset\_1 X3 (k1\_zfmisc\_1 \\ (k3\_cqc\_lang X0))) \Rightarrow ((r3\_cqc\_the1 X0 X3 (k8\_cqc\_lang X0 X1 (k6\_cqc\_lang \\ X0 (k6\_cqc\_lang X0 X2)))) \Leftrightarrow (r3\_cqc\_the1 X0 X3 (k8\_cqc\_lang X0 X1 \\ X2)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m2\_subset\_1 X1 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X2.(m2\_subset\_1 X2 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X3.(m1\_subset\_1 X3 (k1\_zfmisc\_1 \\ (k3\_cqc\_lang X0))) \Rightarrow ((r3\_cqc\_the1 X0 X3 (k8\_cqc\_lang X0 (k6\_cqc\_lang \\ X0 X1) (k6\_cqc\_lang X0 X2))) \Leftrightarrow (r3\_cqc\_the1 X0 X3 (k8\_cqc\_lang X0 \\ X2 X1)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge \\ (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2.(m2\_subset\_1 \\ X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\neg v1\_xboole\_0 (k3\_cqc\_lang X0)) \tag{4}$$

Assume the following.

$$\forall X0.\forall X1.((m1\_qc\_lang1\ X0)\wedge(m1\_subset\_1\ X1\ (k3\_cqc\_lang\ X0)))\Rightarrow(m2\_subset\_1\ (k6\_cqc\_lang\ X0\ X1)\ (k9\_qc\_lang1\ X0)\ (k3\_cqc\_lang\ X0)) \quad (5)$$

Assume the following.

$$\forall X0.(m1\_qc\_lang1\ X0)\Rightarrow(m1\_subset\_1\ (k3\_cqc\_lang\ X0)\ (k1\_zfmisc\_1\ (k9\_qc\_lang1\ X0))) \quad (6)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0\ X0)\Rightarrow(\forall X1.(m1\_subset\_1\ X1\ (k1\_zfmisc\_1\ X0))\Rightarrow(v1\_xboole\_0\ X1)) \quad (7)$$

**Theorem 1**

$$\begin{aligned} \forall X0.(m1\_qc\_lang1\ X0)\Rightarrow(\forall X1.(m2\_subset\_1\ X1\ (k9\_qc\_lang1\ X0)\ (k3\_cqc\_lang\ X0))\Rightarrow(\forall X2.(m2\_subset\_1\ X2\ (k9\_qc\_lang1\ X0)\ (k3\_cqc\_lang\ X0))\Rightarrow(\forall X3.(m1\_subset\_1\ X3\ (k1\_zfmisc\_1\ (k3\_cqc\_lang\ X0))\Rightarrow((r3\_cqc\_the1\ X0\ X3\ (k8\_cqc\_lang\ X0)\ (k6\_cqc\_lang\ X0\ X1)\ X2))\Rightarrow(r3\_cqc\_the1\ X0\ X3\ (k8\_cqc\_lang\ X0)\ (k6\_cqc\_lang\ X0\ X2)\ X1)))))) \end{aligned}$$