

t74_mcart_1

(TMQ_{x2}NXNdUKWc3UaLksoEnVk6Ev3a1a99k)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_xtuple_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_mcart_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_mcart_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_mcart_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_mcart_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k5_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k8_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k7_xtuple_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. \forall X4. ((\neg v1_xboole_0 \\ & X0) \wedge ((\neg v1_xboole_0 X1) \wedge ((\neg v1_xboole_0 X2) \wedge ((\neg v1_xboole_0 X3) \wedge \\ & (m1_subset_1 X4 (k4_zfmisc_1 X0 X1 X2 X3)))))) \Rightarrow (k7_mcart_1 X0 X1 \\ & X2 X3 X4 = k2_xtuple_0 X4) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. \forall X4. ((\neg v1_xboole_0 \\ & X0) \wedge ((\neg v1_xboole_0 X1) \wedge ((\neg v1_xboole_0 X2) \wedge ((\neg v1_xboole_0 X3) \wedge \\ & (m1_subset_1 X4 (k4_zfmisc_1 X0 X1 X2 X3)))))) \Rightarrow (k6_mcart_1 X0 X1 \\ & X2 X3 X4 = k5_xtuple_0 X4) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. \forall X4. ((\neg v1_xboole_0 \\ & X0) \wedge ((\neg v1_xboole_0 X1) \wedge ((\neg v1_xboole_0 X2) \wedge ((\neg v1_xboole_0 X3) \wedge \\ & (m1_subset_1 X4 (k4_zfmisc_1 X0 X1 X2 X3)))))) \Rightarrow (k5_mcart_1 X0 X1 \\ & X2 X3 X4 = k8_xtuple_0 X4) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. \forall X4. ((\neg v1_xboole_0 \\ & X0) \wedge ((\neg v1_xboole_0 X1) \wedge ((\neg v1_xboole_0 X2) \wedge ((\neg v1_xboole_0 X3) \wedge \\ & (m1_subset_1 X4 (k4_zfmisc_1 X0 X1 X2 X3)))))) \Rightarrow (k4_mcart_1 X0 X1 \\ & X2 X3 X4 = k7_xtuple_0 X4) \end{aligned} \tag{4}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.k8_xtuple_0 (k6_xtuple_0 X0 X1 X2 X3) = X1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.k7_xtuple_0 (k6_xtuple_0 X0 X1 X2 X3) = X0 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.k2_xtuple_0 (k6_xtuple_0 X0 X1 X2 X3) = X3 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.k5_xtuple_0 (k6_xtuple_0 X0 X1 X2 X3) = X2 \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(\neg v1_xboole_0 X1) \Rightarrow \\ & (\forall X2.(\neg v1_xboole_0 X2) \Rightarrow (\forall X3.(\neg v1_xboole_0 X3) \Rightarrow \\ & (\forall X4.(m1_subset_1 X4 (k4_zfmisc_1 X0 X1 X2 X3)) \Rightarrow (\forall X5. \\ & \forall X6.\forall X7.\forall X8.(X4 = k6_xtuple_0 X5 X6 X7 X8) \Rightarrow \\ & ((k4_mcart_1 X0 X1 X2 X3 X4 = X5) \wedge ((k5_mcart_1 X0 X1 X2 X3 X4 = X6) \wedge \\ & (k6_mcart_1 X0 X1 X2 X3 X4 = X7) \wedge (k7_mcart_1 X0 X1 X2 X3 X4 = X8)))))))))) \end{aligned}$$