

t74_mesfunc6
(TMbhdCt3UC4LtToWDjYejCk5Kdxk8eqDti9)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k47_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v6_supinf_2 : \iota \Rightarrow o$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k45_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_membered : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge (\\ & \quad m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (\\ & \quad \forall X2. ((v1_funct_1 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & \quad \quad X0 k1_numbers)))) \Rightarrow ((\forall X3. (X3 \in k9_subset_1 X0 (k1_relset_1 \\ & \quad X0 X1) (k1_relset_1 X0 X2)) \Rightarrow (r1_xxreal_0 (k1_seq_1 X2 X3) (k1_seq_1 \\ & \quad \quad X1 X3))) \Rightarrow (v6_supinf_2 (k47_valued_1 X0 k1_numbers k1_numbers \\ & \quad \quad \quad X1 X2)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow \\ & \quad (\forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_valued_0 \\ & \quad X1))) \Rightarrow (k9_xtuple_0 (k45_valued_1 X0 X1) = k3_xboole_0 (k9_xtuple_0 \\ & \quad \quad X0) (k9_xtuple_0 X1))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow (k9_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.\forall X4.((v3_membered X1)\wedge((v3_membered X2)\wedge(((v1_funct_1 X3)\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\wedge((v1_funct_1 X4)\wedge(m1_subset_1 X4 (k1_zfmisc_1 (k2_zfmisc_1 X0 X2)))))))\Rightarrow(k47_valued_1 X0 X1 X2 X3 X4 = k45_valued_1 X3 X4) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (5)$$

Assume the following.

$$v3_membered k1_numbers \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.\forall X4.((v3_membered X1)\wedge((v3_membered X2)\wedge(((v1_funct_1 X3)\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\wedge((v1_funct_1 X4)\wedge(m1_subset_1 X4 (k1_zfmisc_1 (k2_zfmisc_1 X0 X2)))))))\Rightarrow((v1_funct_1 (k47_valued_1 X0 X1 X2 X3 X4)\wedge(m1_subset_1 (k47_valued_1 X0 X1 X2 X3 X4) (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_valued_0 X0)))\wedge((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_valued_0 X1))))\Rightarrow((v1_relat_1 (k45_valued_1 X0 X1))\wedge(v1_funct_1 (k45_valued_1 X0 X1))) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(m1_subset_1 (k1_relset_1 X0 X1) (k1_zfmisc_1 X0)) \quad (9)$$

Assume the following.

$$\forall X0.(v3_membered X0)\Rightarrow(v1_membered X0) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.(v1_membered\ X1)\Rightarrow(\forall X2.(m1_subset_1\ X2\ (k1_zfmisc_1\ (k2_zfmisc_1\ X0\ X1)))\Rightarrow(v1_valued_0\ X2)) \quad (13)$$

Theorem 1

$$\begin{aligned} \forall X0.(\neg v1_xboole_0\ X0)\Rightarrow(\forall X1.((v1_funct_1\ X1)\wedge(\\ m1_subset_1\ X1\ (k1_zfmisc_1\ (k2_zfmisc_1\ X0\ k1_numbers))))\Rightarrow(\\ \forall X2.((v1_funct_1\ X2)\wedge(m1_subset_1\ X2\ (k1_zfmisc_1\ (k2_zfmisc_1 \\ X0\ k1_numbers))))\Rightarrow((\forall X3.(X3 \in k1_relset_1\ X0\ (k47_valued_1 \\ X0\ k1_numbers\ k1_numbers\ X1\ X2))\Rightarrow(r1_xxreal_0\ (k1_seq_1\ X2\ X3) \\ (k1_seq_1\ X1\ X3)))\Rightarrow(v6_supinf_2\ (k47_valued_1\ X0\ k1_numbers\ k1_numbers \\ X1\ X2)))))) \end{aligned}$$