

t74\_relat\_1 (TM-  
bxBkyTG5EJVMiXjV3QFK5gXpaf8VMQVpx)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (v1\_relat\_1 X2) \Rightarrow (k5\_relat\_1 (k5\_relat\_1 X2 X0) X1 = k5\_relat\_1 X2 (k3\_xboole\_0 X0 X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (r1\_tarski X0 X1) \Rightarrow (k3\_xboole\_0 X0 X1 = X0) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. k3\_xboole\_0 X0 X1 = k3\_xboole\_0 X1 X0 \quad (3)$$

**Theorem 1**

$$\forall X0. \forall X1. \forall X2. (v1\_relat\_1 X2) \Rightarrow ((r1\_tarski X0 X1) \Rightarrow (k5\_relat\_1 (k5\_relat\_1 X2 X1) X0 = k5\_relat\_1 X2 X0))$$