

t75_finseq_5

(TMH49STGbLhS5eyMBWaPTcG96E7oB7UuU7D)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_finseq_5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k17_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k16_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(\neg v1_xboole_0 X1) \Rightarrow (\\ \forall X2.(m1_subset_1 X2 X1) \Rightarrow (\forall X3.(m2_finseq_1 X3 X1) \Rightarrow \\ (\forall X4.(v7_ordinal1 X4) \Rightarrow ((X4 \in k4_finseq_1 (k17_finseq_1 \\ X1 X0 X3)) \Rightarrow (k7_partfun1 X1 (k5_finseq_5 X1 X3 X2 X0) X4 = k7_partfun1 \\ X1 X3 X4)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge ((\\ v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow ((r1_xxreal_0 X0 (k3_finseq_1 \\ X1) \Rightarrow (k3_finseq_1 (k16_finseq_1 X0 X1) = X0))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge ((\\ v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow ((r1_xxreal_0 (k3_finseq_1 \\ X1) X0) \Rightarrow (k16_finseq_1 X0 X1 = X1))) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ (\forall X1.(v7_ordinal1 X1) \Rightarrow ((X1 \in k1_relset_1 k5_numbers X0) \Leftrightarrow \\ ((r1_xxreal_0 np_1 X1) \wedge (r1_xxreal_0 X1 (k3_finseq_1 X0)))))) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow ((\neg r1_xxreal_0 np_1 X0) \Rightarrow (X0 = k6_numbers)) \quad (5)$$

Assume the following.

$$\begin{aligned} ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (6)$$

Assume the following.

$$r1_xxreal_0 np_1 np_1 \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (9)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (10)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ (k4_finseq_1 X0 = k9_xtuple_0 X0) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (\\ k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v7_ordinal1 X1) \wedge (m1_finseq_1 \\ X2 X0)) \Rightarrow (k17_finseq_1 X0 X1 X2 = k16_finseq_1 X1 X2) \quad (13)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (14)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & ((v1_relat_1 (k16_finseq_1 k6_numbers X0)) \wedge ((v1_funct_1 (k16_finseq_1 \\ & k6_numbers X0)) \wedge ((v1_xboole_0 (k16_finseq_1 k6_numbers X0)) \wedge \\ & (v1_finseq_1 (k16_finseq_1 k6_numbers X0)))))) \end{aligned} \quad (15)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m2_finseq_1 X1 X0) \Rightarrow ((v1_funct_1 X1) \wedge (\\ & (v1_finseq_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers \\ & X0)))))) \end{aligned} \quad (16)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_finseq_1 X1 X0) \Rightarrow ((v1_relat_1 X1) \wedge (\\ & (v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \end{aligned} \quad (17)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (18)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (m2_subset_1 (k3_finseq_1 X0) k1_numbers k5_numbers) \end{aligned} \quad (19)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v7_ordinal1 X0) \wedge ((v1_relat_1 X1) \wedge (\\ & v1_funct_1 X1) \wedge (v1_finseq_1 X1)))) \Rightarrow ((v1_relat_1 (k16_finseq_1 \\ & X0 X1)) \wedge ((v1_funct_1 (k16_finseq_1 X0 X1)) \wedge (v1_finseq_1 (k16_finseq_1 \\ & X0 X1)))) \end{aligned} \quad (20)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (\\ & (r1_xxreal_0 X0 X1) \vee (r1_xxreal_0 X1 X0)) \end{aligned} \quad (21)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (22)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge \\ & (v1_finseq_1 X0)))) \end{aligned} \quad (23)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(v1_xreal_0\ X0) \quad (24)$$

Assume the following.

$$\forall X0.(v1_xboole_0\ X0)\Rightarrow(\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1\ X0))\Rightarrow(v1_xboole_0\ X1)) \quad (25)$$

Theorem 1

$$\begin{aligned} &\forall X0.(v7_ordinal1\ X0)\Rightarrow(\forall X1.(\neg v1_xboole_0\ X1)\Rightarrow(\\ &\quad \forall X2.(m1_subset_1\ X2\ X1)\Rightarrow(\forall X3.(m2_finseq_1\ X3\ X1)\Rightarrow \\ &((r1_xreal_0\ np_1\ X0)\Rightarrow((v1_xboole_0\ X3)\vee(k7_partfun1\ X1\ (k5_finseq_5 \\ &\quad X1\ X3\ X2\ X0)\ np_1 = k7_partfun1\ X1\ X3\ np_1)))))) \end{aligned}$$