

t75_funcop_1
(TMaF7tz3JVhEZMTRaMjZbWkj7Ku7Jfy7NHx)

October 27, 2020

Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k16_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k7_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (k9_xtuple_0 (k2_funcop_1 X0 X1) = X0) \wedge (r1_tarski (k10_xtuple_0 (k2_funcop_1 X0 X1)) (k1_tarski X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. k7_funcop_1 X0 X1 = k2_funcop_1 X0 X1 \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. k16_funcop_1 X0 X1 = k7_funcop_1 (k1_tarski X0) X1 \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (4)$$

Theorem 1

$$\forall X0. \forall X1. \forall X2. (X0 \in k9_xtuple_0 (k16_funcop_1 X1 X2)) \Rightarrow (X0 = X1)$$