

t75\_funct\_2  
(TMUUHQ11Gx1tkdy1tb6XUvBDJTLHv7LD5P3)

October 27, 2020

Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $r1\_partfun1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $r1\_tarSKI : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.(( \\ v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((r1\_tarSKI (k9\_xtuple\_0 X0) \\ (k9\_xtuple\_0 X1)) \Rightarrow ((r1\_partfun1 X0 X1) \Leftrightarrow (\forall X2.(X2 \in k9\_xtuple\_0 \\ X0) \Rightarrow (k1\_funct\_1 X0 X2 = k1\_funct\_1 X1 X2)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_subset\_1 X0 (k1\_zfmisc\_1 X1)) \Leftrightarrow (r1\_tarSKI X0 X1) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \wedge (( \\ v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1))) \Rightarrow (r1\_partfun1 X0 X0) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow ( \\ k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (5)$$

Assume the following.

$$v1\_xboole\_0 k1\_xboole\_0 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1)\wedge(v4\_relat\_1 X1 X0))\Rightarrow(m1\_subset\_1 (k1\_relset\_1 X0 X1) (k1\_zfmisc\_1 X0)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(((X1\neq k1\_xboole\_0)\Rightarrow((v1\_funct\_2 X2 X0 X1)\Leftrightarrow(X0 = k1\_relset\_1 X0 X2)))\wedge((X1 = k1\_xboole\_0)\Rightarrow((v1\_funct\_2 X2 X0 X1)\Leftrightarrow(X2 = k1\_xboole\_0)))) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(v1\_xboole\_0 X0)\Rightarrow(\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X1 X0)))\Rightarrow(v1\_xboole\_0 X2)) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow((v4\_relat\_1 X2 X0)\wedge(v5\_relat\_1 X2 X1)) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(v1\_relat\_1 X2) \quad (11)$$

**Theorem 1**

$$\forall X0.\forall X1.\forall X2.((v1\_funct\_1 X2)\wedge(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))))\Rightarrow(\forall X3.((v1\_funct\_1 X3)\wedge((v1\_funct\_2 X3 X0 X1)\wedge(m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))))))\Rightarrow(((X1 = k1\_xboole\_0)\Rightarrow(X0 = k1\_xboole\_0))\Rightarrow((r1\_partfun1 X2 X3)\Leftrightarrow(\forall X4.(X4 \in k1\_relset\_1 X0 X2)\Rightarrow(k1\_funct\_1 X2 X4 = k1\_funct\_1 X3 X4))))$$