

t75_matrixr2 (TMTGe- wax3avc4DGmSNYfwhc7LzK2LfmYjXS)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_matrixr2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_funct_7 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k5_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_card_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. \neg (X0 \in X1) \wedge ((m1_subset_1 X1 (k1_zfmisc_1 X2)) \wedge (v1_xboole_0 X2)) \quad (1)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. (X2 \in k9_xtuple_0 X0) \Rightarrow (k1_funct_1 (k2_funct_7 X0 X2 X1) X2 = X1)) \quad (2)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (\forall X1. (v7_ordinal1 X1) \Rightarrow ((X1 \in k1_relset_1 k5_numbers X0) \Leftrightarrow ((r1_xxreal_0 np_1 X1) \wedge (r1_xxreal_0 X1 (k3_finseq_1 X0)))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((\neg v1_xboole_0 X0) \wedge ((v7_ordinal1 X1) \wedge (m1_subset_1 X2 X0))) \Rightarrow (k5_finseq_2 X0 X1 X2 = k2_finseq_2 X1 X2) \quad (4)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k3_finseq_1 X0 = k1_card_1 X0) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(v7_ordinal1 X0)\Rightarrow((v1_relat_1 (k2_finseq_2 X0 X1))\wedge((v1_funct_1 (k2_finseq_2 X0 X1))\wedge((v3_card_1 (k2_finseq_2 X0 X1) X0)\wedge(v1_finseq_1 (k2_finseq_2 X0 X1)))))) \quad (7)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((\neg v1_xboole_0 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 X0))))\Rightarrow(\forall X2.(m2_subset_1 X2 X0 X1)\Rightarrow(m1_subset_1 X2 X0)) \quad (9)$$

Assume the following.

$$m2_subset_1 k6_numbers k1_numbers k5_numbers \quad (10)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.(v7_ordinal1 X0)\Rightarrow((v1_relat_1 (k2_finseq_2 X0 X1))\wedge((v1_funct_1 (k2_finseq_2 X0 X1))\wedge(v1_finseq_1 (k2_finseq_2 X0 X1)))) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(m1_subset_1 (k1_relset_1 X0 X1) (k1_zfmisc_1 X0)) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.(v3_card_1 X1 X0)\Leftrightarrow(k1_card_1 X1 = X0) \quad (14)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(\forall X1.(v7_ordinal1 X1)\Rightarrow(k6_matrixr2 X0 X1 = k2_funct_7 (k5_finseq_2 k1_numbers X0 k6_numbers) X1 np_1)) \quad (15)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow((v1_relat_1 X0)\wedge((v4_relat_1 X0 k5_numbers)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))) \quad (16)$$

Theorem 1

$$\forall X0.(v7_ordinal1\ X0) \Rightarrow (\forall X1.(v7_ordinal1\ X1) \Rightarrow ((r1_xxreal_0\ np_1\ X0) \wedge (r1_xxreal_0\ X0\ X1)) \Rightarrow (k1_funct_1\ (k6_matrixr2\ X1\ X0)\ X0 = np_1)))$$