

t75_sin_cos6 (TM-
daY1z5tAXigceLrW5VQUAEqNZivMpqFTz)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_real_1 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k3_sin_cos6 : \iota \Rightarrow \iota$ be given. Let $k10_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k32_sin_cos : \iota$ be given. Let $np_2 : \iota$ be given. Let $k21_sin_cos : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k18_sin_cos : \iota \Rightarrow \iota$ be given. Let $k7_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned}
 & (k21_sin_cos (k10_real_1 k32_sin_cos np_2) = k6_numbers) \wedge ((\\
 & k18_sin_cos (k10_real_1 k32_sin_cos np_2) = np_1) \wedge ((k21_sin_cos \\
 & k32_sin_cos = k1_real_1 np_1) \wedge ((k18_sin_cos k32_sin_cos = k6_numbers) \wedge \\
 & ((k21_sin_cos (k7_real_1 k32_sin_cos (k10_real_1 k32_sin_cos \\
 & np_2)) = k6_numbers) \wedge ((k18_sin_cos (k7_real_1 k32_sin_cos (\\
 & k10_real_1 k32_sin_cos np_2)) = k1_real_1 np_1) \wedge ((k21_sin_cos \\
 & (k8_real_1 np_2 k32_sin_cos) = np_1) \wedge (k18_sin_cos (k8_real_1 \\
 & np_2 k32_sin_cos) = k6_numbers))))))
 \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. (v1_xreal_0 X0) \Rightarrow (((r1_xxreal_0 (k1_real_1 np_1) X0) \wedge (r1_xxreal_0 X0 np_1)) \Rightarrow (k18_sin_cos (k3_sin_cos6 X0) = X0)) \tag{2}$$

Theorem 1

$$\forall X0. (v1_xreal_0 X0) \Rightarrow (((r1_xxreal_0 (k1_real_1 np_1) X0) \wedge ((r1_xxreal_0 X0 np_1) \wedge (k3_sin_cos6 X0 = k10_real_1 k32_sin_cos np_2))) \Rightarrow (X0 = np_1))$$