

t75_xxreal_1

(TMZighmM8P9eRP1upcMQtqnqGZfBqJSG4vo)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3.(v1_xxreal_0 X3) \Rightarrow ((r1_tarski (\\ & k2_xxreal_1 X0 X1) (k4_xxreal_1 X2 X3)) \Rightarrow ((r1_xxreal_0 X1 X0) \vee (\\ & (\neg r1_xxreal_0 X0 X2) \wedge (r1_xxreal_0 X1 X3))))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3.(v1_xxreal_0 X3) \Rightarrow ((r1_tarski (\\ & k4_xxreal_1 X0 X1) (k2_xxreal_1 X2 X3)) \Rightarrow ((r1_xxreal_0 X1 X0) \vee (\\ & (r1_xxreal_0 X2 X0) \wedge (r1_xxreal_0 X1 X3))))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow \\ & (r1_xxreal_0 X0 X2)))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. r1_tarski X0 X0 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (\\ (r1_xxreal_0 X0 X1) \vee (r1_xxreal_0 X1 X0)) \quad (5)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3.(v1_xxreal_0 X3) \Rightarrow (\neg(\neg r1_xxreal_0 \\ & X1 X0) \wedge (k2_xxreal_1 X0 X1 = k4_xxreal_1 X2 X3)))))) \end{aligned}$$