

## t76\_finseq\_6

(TMGimv4iKbx8tz4eUsJSfUSV22k45igUsdy)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k8\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_finseq\_5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_rfinseq : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k2\_finseq\_5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k7\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_finseq\_4 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k6\_finseq\_4 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k12\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_numbers : \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ & (\forall X1.(X1 \in k10\_xtuple\_0 X0) \Rightarrow (X0 = k7\_finseq\_1 (k7\_finseq\_1 \\ & (k5\_finseq\_4 X0 X1) (k9\_finseq\_1 X1)) (k6\_finseq\_4 X0 X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 X0) \Rightarrow \\ & (\forall X2.(m2\_finseq\_1 X2 X0) \Rightarrow (k2\_rfinseq X0 np\_1 (k8\_finseq\_1 \\ & X0 (k12\_finseq\_1 X0 X1) X2) = X2))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ & (\forall X1.(X1 \in k10\_xtuple\_0 X0) \Rightarrow (r1\_tarski (k10\_xtuple\_0 ( \\ & k6\_finseq\_4 X0 X1)) (k10\_xtuple\_0 X0))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 X0) \Rightarrow \\ & (\forall X2.(m2\_finseq\_1 X2 X0) \Rightarrow ((X1 \in k10\_xtuple\_0 X2) \Rightarrow (k2\_finseq\_5 \\ & X0 X2 X1 = k7\_finseq\_1 (k12\_finseq\_1 X0 X1) (k6\_finseq\_4 X2 X1)))))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 X0) \Rightarrow \\ & (\forall X2.(m2\_finseq\_1 X2 X0) \Rightarrow ((X1 \in k10\_xtuple\_0 X2) \Rightarrow (k1\_finseq\_5 \\ & X0 X2 X1 = k7\_finseq\_1 (k5\_finseq\_4 X2 X1) (k12\_finseq\_1 X0 X1)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((r1\_tarski X0 X1) \wedge (r1\_tarski X1 X2)) \Rightarrow (r1\_tarski X0 X2) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0) \Leftrightarrow (m1\_finseq\_1 X1 X0) \quad (7)$$

Assume the following.

$$\forall X0.k9\_finseq\_1 X0 = k5\_finseq\_1 X0 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1\_finseq\_1 X1 X0) \wedge (m1\_finseq\_1 X2 X0)) \Rightarrow (k8\_finseq\_1 X0 X1 X2 = k7\_finseq\_1 X1 X2) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0) \wedge (m1\_subset\_1 X1 X0)) \Rightarrow (k12\_finseq\_1 X0 X1 = k5\_finseq\_1 X1) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0) \Rightarrow ((v1\_funct\_1 X1) \wedge ((v1\_finseq\_1 X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers X0)))))) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_1 X1 X0) \Rightarrow ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow ((v1\_relat\_1 (k6\_finseq\_4 X0 X1)) \wedge ((v1\_funct\_1 (k6\_finseq\_4 X0 X1)) \wedge (v1\_finseq\_1 (k6\_finseq\_4 X0 X1)))) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1\_xboole\_0 X0) \wedge (m1\_finseq\_1 X1 X0)) \Rightarrow (m2\_finseq\_1 (k1\_finseq\_5 X0 X1 X2) X0) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0)\wedge(m1\_subset\_1 X1 X0))\Rightarrow (m2\_finseq\_1 (k12\_finseq\_1 X0 X1) X0) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1)\wedge((v1\_funct\_1 X1)\wedge(v1\_finseq\_1 X1)))\Rightarrow((m1\_finseq\_1 X1 X0)\Leftrightarrow(r1\_tarski (k10\_xtuple\_0 X1) X0)) \quad (16)$$

**Theorem 1**

$$\begin{aligned} &\forall X0.(\neg v1\_xboole\_0 X0)\Rightarrow(\forall X1.(m1\_subset\_1 X1 X0)\Rightarrow \\ &(\forall X2.(m2\_finseq\_1 X2 X0)\Rightarrow((X1 \in k10\_xtuple\_0 X2)\Rightarrow(k8\_finseq\_1 \\ &X0 (k1\_finseq\_5 X0 X2 X1) (k2\_rfinseq X0 np\_1 (k2\_finseq\_5 X0 X2 \\ &X1) = X2)))) \end{aligned}$$