

t76_funct_7 (TMYNGN- DAoX87uaaT8hiKytp2kMXx2ktC6br)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k3_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_funct_7 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_partfun1 : \iota \Rightarrow \iota$ be given. Let $k1_relat_1 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k4_relat_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(m1_subset_1 X0 k5_numbers) \Rightarrow (\forall X1.(v1_relat_1 X1) \Rightarrow (k3_relat_1 (k6_partfun1 (k1_relat_1 X1)) (k9_funct_7 X1 X0) = k9_funct_7 X1 X0)) \quad (1)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k5_numbers) \Rightarrow (\forall X1.(v1_relat_1 X1) \Rightarrow ((r1_tarski (k9_xtuple_0 (k9_funct_7 X1 X0)) (k1_relat_1 X1)) \wedge (r1_tarski (k10_xtuple_0 (k9_funct_7 X1 X0)) (k1_relat_1 X1)))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(v1_relat_1 X1) \Rightarrow ((r1_tarski (k10_xtuple_0 X1) X0) \Rightarrow (k3_relat_1 X1 (k4_relat_1 X0) = X1)) \quad (3)$$

Assume the following.

$$\forall X0.k6_partfun1 X0 = k4_relat_1 X0 \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.v1_relat_1 (k3_relat_1 X0 X1) \quad (5)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 k5_numbers) \Rightarrow (\forall X1.(v1_relat_1 X1) \Rightarrow (k3_relat_1 (k9_funct_7 X1 X0) (k6_partfun1 (k1_relat_1 X1)) = k9_funct_7 X1 X0))$$