

t76_pdiff_5

(TMV5okmUUGAodwuKwMYwjSCuMK5dHhxsTze)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_euclid : \iota \Rightarrow \iota$ be given. Let $np_3 : \iota$ be given. Let $k1_numbers : \iota$ be given. Let $m2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_pdiff_5 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_fcont_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_pdiff_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_pdiff_3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_pdiff_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r3_pdiff_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v2_xreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0. ((v1_funct_1 X0) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \\ (k1_euclid np_3) k1_numbers)))) \Rightarrow (\forall X1. (m2_finseq_2 X1 \\ k1_numbers (k1_euclid np_3)) \Rightarrow ((r3_pdiff_1 np_3 np_1 X0 X1) \Rightarrow \\ (r1_fcont_1 (k1_pdiff_2 np_3 np_1 X0 X1) (k1_seq_1 (k1_pdiff_1 \\ np_1 np_3) X1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. (m2_finseq_2 X0 k1_numbers (k1_euclid np_3)) \Rightarrow (\forall X1. \\ ((v1_funct_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 (\\ k1_euclid np_3) k1_numbers)))) \Rightarrow ((r1_pdiff_5 X1 X0) \Leftrightarrow (r3_pdiff_1 \\ np_3 np_1 (k1_pdiff_3 np_1 np_3 X1) X0))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} ((v2_xreal_0 np_3) \wedge (m2_subset_1 np_3 k1_numbers k5_numbers)) \wedge \\ ((m1_subset_1 np_3 k5_numbers) \wedge (m1_subset_1 np_3 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\neg v1_xboole_0 np_3 \quad (4)$$

Assume the following.

$$\begin{aligned} ((v2_xreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned}
& \forall X0.\forall X1.\forall X2.((m1_subset_1 X0 k5_numbers)\wedge \\
& (((\neg v1_xboole_0 X1)\wedge(m1_subset_1 X1 k5_numbers))\wedge((v1_funct_1 \\
& X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 (k1_euclid X1) \\
& k1_numbers))))))\Rightarrow((v1_funct_1 (k1_pdiff_3 X0 X1 X2))\wedge((v1_funct_2 \\
& (k1_pdiff_3 X0 X1 X2) (k1_euclid X1) k1_numbers)\wedge(m1_subset_1 \\
& (k1_pdiff_3 X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 (k1_euclid X1) \\
& k1_numbers))))))
\end{aligned} \tag{6}$$

Theorem 1

$$\begin{aligned}
& \forall X0.((v1_funct_1 X0)\wedge(m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \\
& (k1_euclid np_3) k1_numbers))))\Rightarrow(\forall X1.(m2_finseq_2 X1 \\
& k1_numbers (k1_euclid np_3))\Rightarrow((r1_pdiff_5 X0 X1)\Rightarrow(r1_fcont_1 \\
& (k1_pdiff_2 np_3 np_1 (k1_pdiff_3 np_1 np_3 X0) X1) (k1_seq_1 \\
& (k1_pdiff_1 np_1 np_3) X1))))
\end{aligned}$$