

t76_sheffer2

(TMK_m43qv5Qx6chwvf76MLjoUFdrW34gVuye)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v10_sheffer1 : \iota \Rightarrow o$ be given. Let $v11_sheffer1 : \iota \Rightarrow o$ be given. Let $v12_sheffer1 : \iota \Rightarrow o$ be given. Let $l1_sheffer1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k5_sheffer1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v10_sheffer1 X0) \wedge ((v11_sheffer1 \\ & X0) \wedge ((v12_sheffer1 X0) \wedge (l1_sheffer1 X0)))))) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (u1_struct_0 \\ & X0)) \Rightarrow (k5_sheffer1 X0 X2 (k5_sheffer1 X0 (k5_sheffer1 X0 X1 X1) X1) = \\ & k5_sheffer1 X0 X2 X2))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(((\neg v2_struct_0 X0) \wedge (l1_sheffer1 \\ & X0)) \wedge ((m1_subset_1 X1 (u1_struct_0 X0)) \wedge (m1_subset_1 X2 (u1_struct_0 \\ & X0)))) \Rightarrow (m1_subset_1 (k5_sheffer1 X0 X1 X2) (u1_struct_0 X0)) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (l1_sheffer1 X0)) \Rightarrow ((v10_sheffer1 \\ & X0) \Leftrightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (k5_sheffer1 \\ & X0 (k5_sheffer1 X0 X1 X1) (k5_sheffer1 X0 X1 X1) = X1))) \end{aligned} \tag{3}$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v10_sheffer1 X0) \wedge ((v11_sheffer1 \\ & X0) \wedge ((v12_sheffer1 X0) \wedge (l1_sheffer1 X0)))))) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (u1_struct_0 \\ & X0)) \Rightarrow (X2 = k5_sheffer1 X0 (k5_sheffer1 X0 X2 X2) (k5_sheffer1 X0 \\ & (k5_sheffer1 X0 X1 X1) X1)))) \end{aligned}$$