

t77_pzfmisc1

(TMRcWzi3f3YZek92ijMtL7WSrSFW98ridLf)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r2_pboole : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_pboole : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v4_relat_1 X1 X0) \wedge \\ & (v1_funct_1 X1) \wedge (v1_partfun1 X1 X0))) \Rightarrow (\forall X2. ((v1_relat_1 \\ & X2) \wedge ((v4_relat_1 X2 X0) \wedge ((v1_funct_1 X2) \wedge (v1_partfun1 X2 X0)))) \Rightarrow \\ & (\forall X3. ((v1_relat_1 X3) \wedge ((v4_relat_1 X3 X0) \wedge ((v1_funct_1 \\ & X3) \wedge (v1_partfun1 X3 X0)))) \Rightarrow (\forall X4. ((v1_relat_1 X4) \wedge ((v4_relat_1 \\ & X4 X0) \wedge ((v1_funct_1 X4) \wedge (v1_partfun1 X4 X0)))) \Rightarrow (((r2_pboole \\ & X0 X1 X2) \wedge (r2_pboole X0 X3 X4)) \Rightarrow (r2_pboole X0 (k6_pboole X0 X1 X3) \\ & (k6_pboole X0 X2 X4)))))) \end{aligned} \tag{1}$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v4_relat_1 X1 X0) \wedge \\ & (v1_funct_1 X1) \wedge (v1_partfun1 X1 X0))) \Rightarrow (\forall X2. ((v1_relat_1 \\ & X2) \wedge ((v4_relat_1 X2 X0) \wedge ((v1_funct_1 X2) \wedge (v1_partfun1 X2 X0)))) \Rightarrow \\ & ((r2_pboole X0 X1 X2) \Rightarrow (r2_pboole X0 (k6_pboole X0 X1 X1) (k6_pboole \\ & X0 X2 X2)))) \end{aligned}$$