

t77_xboole_1

(TMYLMwAf6VVm4h21wL99R2UsgLiRSJ9wkxC)

October 27, 2020

Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Rightarrow (k3_xboole_0 X0 X1 = X0) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. k3_xboole_0 (k3_xboole_0 X0 X1) X2 = k3_xboole_0 X0 (k3_xboole_0 X1 X2) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1_xboole_0 X0 X1) \Leftrightarrow (k3_xboole_0 X0 X1 = k1_xboole_0) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (4)$$

Theorem 1

$$\forall X0. \forall X1. \forall X2. \neg(\neg r1_xboole_0 X0 X1) \wedge ((r1_tarski X0 X2) \wedge (r1_xboole_0 X0 (k3_xboole_0 X1 X2)))$$