

t78_funct_1
(TMJzu51DS1mLcmudMsKuKxvjsx9r9zqt8MRU)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_tarski X0 (k10_xtuple_0 X1)) \Rightarrow (k7_relat_1 X1 (k8_relat_1 X1 X0) = X0)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. r1_tarski (k3_xboole_0 X0 X1) X0 \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (k8_relat_1 X1 X0 = k8_relat_1 X1 (k3_xboole_0 (k10_xtuple_0 X1) X0)) \quad (3)$$

Assume the following.

$$\forall X0. (v1_relat_1 X0) \Rightarrow (k7_relat_1 X0 (k9_xtuple_0 X0) = k10_xtuple_0 X0) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (5)$$

Theorem 1

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (k7_relat_1 X1 (k8_relat_1 X1 X0) = k3_xboole_0 X0 (k7_relat_1 X1 (k9_xtuple_0 X1)))$$