

## t78\_setlim\_1

(TMHsov5FWwCviVCwknD5rM6fBRBkkN7PspT)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_prob\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v4\_prob\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k9\_setfam\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v3\_prob\_1 : \iota \Rightarrow o$  be given. Let  $k4\_kurato\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_kurato\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v3\_kurato\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_funct\_1 X1) \wedge ((v1\_funct\_2 X1 k5\_numbers \\ & (k9\_setfam\_1 X0)) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & k5\_numbers (k9\_setfam\_1 X0)))))) \Rightarrow ((v3\_prob\_1 X1) \Rightarrow ((v3\_kurato\_0 \\ & X1 X0) \wedge (k4\_kurato\_0 X0 X1 = k1\_kurato\_0 X0 X1))) \end{aligned} \quad (1)$$

### Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1\_xboole\_0 X1) \wedge ((v1\_prob\_1 X1 X0) \wedge \\ & ((v4\_prob\_1 X1 X0) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k1\_zfmisc\_1 \\ & X0)))))) \Rightarrow (\forall X2. ((v5\_relat\_1 X2 X1) \wedge ((v1\_funct\_1 X2) \wedge \\ & (v1\_funct\_2 X2 k5\_numbers (k9\_setfam\_1 X0)) \wedge (m1\_subset\_1 X2 ( \\ & k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers (k9\_setfam\_1 X0)))))) \Rightarrow \\ & ((v3\_prob\_1 X2) \Rightarrow (k4\_kurato\_0 X0 X2 = k1\_kurato\_0 X0 X2))) \end{aligned}$$