

t78_zf_lang1

(TMXZx3RLVwew4PkrF5SwWYdBKQHhvwDaxUL)

October 27, 2020

Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k1_zf_lang : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_zf_model : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_zf_lang : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge (\\ & (v1_funct_2 X1 k1_zf_lang X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k1_zf_lang X0)))))) \Rightarrow (\forall X2. (m2_subset_1 X2 k5_numbers k1_zf_lang) \Rightarrow \\ & (\forall X3. (m2_subset_1 X3 k5_numbers k1_zf_lang) \Rightarrow ((r1_zf_model \\ & X0 X1 (k4_zf_lang X2 X3)) \Leftrightarrow (k3_funct_2 k1_zf_lang X0 X1 X2 = k3_funct_2 \\ & k1_zf_lang X0 X1 X3)))))) \end{aligned} \tag{1}$$

Theorem 1

$$\begin{aligned} & \forall X0. (m2_subset_1 X0 k5_numbers k1_zf_lang) \Rightarrow (\forall X1. \\ & (\neg v1_xboole_0 X1) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge ((v1_funct_2 \\ & X2 k1_zf_lang X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 k1_zf_lang \\ & X1)))))) \Rightarrow (r1_zf_model X1 X2 (k4_zf_lang X0 X0)))) \end{aligned}$$