

t78\_zfmisc\_1 (TMa-  
JqpCkCd5G4eHdmHsBzw43JVqEmXEdC9z)

October 27, 2020

Let  $k3\_tarski : \iota \Rightarrow \iota$  be given. Let  $k2\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. r1\_tarski X0 (k2\_xboole\_0 X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k3\_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. (X2 \in X3) \wedge (X3 \in X0))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k2\_xboole\_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \vee (X3 \in X1))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (r1\_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. k2\_xboole\_0 X0 X1 = k2\_xboole\_0 X1 X0 \quad (5)$$

**Theorem 1**

$$\forall X0. \forall X1. k3\_tarski (k2\_xboole\_0 X0 X1) = k2\_xboole\_0 (k3\_tarski X0) (k3\_tarski X1)$$