

t79_euclid.8 (TMNRELDVtXtc- tQg1cUQMwCt6caBHgg3TToA)

October 27, 2020

Let $m2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k1_euclid : \iota \Rightarrow \iota$ be given. Let $np_3 : \iota$ be given. Let $k23_rvsum_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k16_euclid : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k5_euclid : \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k4_euclid : \iota \Rightarrow \iota$ be given. Let $c2_topalg_3 : \iota$ be given. Assume the following.

$$\forall X0.(m2_finseq_2 X0 k1_numbers (k1_euclid np_3)) \Rightarrow (k23_rvsum_1 X0 (k16_euclid np_3) = k6_numbers) \quad (1)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(m2_finseq_2 X1 k1_numbers (k1_euclid X0)) \Rightarrow ((k23_rvsum_1 X1 X1 = k6_numbers) \Leftrightarrow (X1 = k5_euclid X0))) \quad (2)$$

Assume the following.

$$((v2_xxreal_0 np_3) \wedge (m2_subset_1 np_3 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_3 k5_numbers) \wedge (m1_subset_1 np_3 k1_numbers)) \quad (3)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (4)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (k5_euclid X0 = k4_euclid X0) \quad (6)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(k16_euclid\ X0 = k4_euclid\ X0) \quad (7)$$

Assume the following.

$$c2_topalg_3 = k6_numbers \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k4_ordinal1)\Rightarrow(v7_ordinal1\ X0) \quad (9)$$

Theorem 1

$$\forall X0.(m2_finseq_2\ X0\ k1_numbers\ (k1_euclid\ np_3))\Rightarrow((k23_rvsum_1\ X0\ X0 = k6_numbers)\Leftrightarrow(X0 = k16_euclid\ np_3))$$