

t79_flang_3
(TMU1SHkBQ4guPmSw8Poi8uiojksRt98WpMN)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k8_afinsq_1 : \iota \Rightarrow \iota$ be given. Let $k6_flang_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_flang_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_flang_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_flang_3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 X0))) \Rightarrow (k2_flang_3 X0 X1 = k1_flang_3 X0 X1 np_1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 X0))) \Rightarrow (\forall X2. (v7_ordinal1 X2) \Rightarrow (k6_flang_1 X0 (k1_flang_3 X0 X1 X2) (k2_flang_2 X0 X1) = k6_flang_1 X0 (k2_flang_2 X0 X1) (k1_flang_3 X0 X1 X2))) \quad (2)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (3)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (5)$$

Theorem 1

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 X0))) \Rightarrow (k6_flang_1 X0 (k2_flang_3 X0 X1) (k2_flang_2 X0 X1) = k6_flang_1 X0 (k2_flang_2 X0 X1) (k2_flang_3 X0 X1))$$