

t79\_glib\_001

(TMKwz6zRnx1zx1nVSk6qpq82cqXA9tQeqKG)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finset\_1 : \iota \Rightarrow o$  be given. Let  $v1\_glib\_000 : \iota \Rightarrow o$  be given. Let  $m3\_glib\_001 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k12\_glib\_001 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_glib\_000 : \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $m2\_glib\_001 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((X0 \in k9\_xtuple\_0 X1) \Rightarrow (k1\_funct\_1 X1 X0 \in k10\_xtuple\_0 X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m2\_finseq\_1 X1 X0) \Leftrightarrow (m1\_finseq\_1 X1 X0) \quad (2)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (4)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v4\_relat\_1 X0 k5\_numbers) \wedge ((v1\_funct\_1 X0) \wedge ((v1\_finset\_1 X0) \wedge (v1\_glib\_000 X0))))) \Rightarrow (\forall X1. (m2\_glib\_001 X1 X0) \Rightarrow (m2\_finseq\_1 X1 (k7\_glib\_000 X0))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0)\Rightarrow((v1\_funct\_1 X1)\wedge((v1\_finseq\_1 X1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers X0)))))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_1 X1 X0)\Rightarrow((v1\_relat\_1 X1)\wedge((v1\_funct\_1 X1)\wedge(v1\_finseq\_1 X1))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(((v1\_relat\_1 X0)\wedge((v4\_relat\_1 X0 k5\_numbers)\wedge((v1\_funct\_1 X0)\wedge((v1\_finset\_1 X0)\wedge(v1\_glib\_000 X0))))))\wedge(m3\_glib\_001 X1 X0)\Rightarrow(m2\_glib\_001 (k12\_glib\_001 X0 X1) X0) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(r1\_tarski X0 X1)\Leftrightarrow(\forall X2.(X2 \in X0)\Rightarrow(X2 \in X1)) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(v1\_relat\_1 X1)\Rightarrow((v5\_relat\_1 X1 X0)\Leftrightarrow(r1\_tarski (k10\_xtuple\_0 X1) X0)) \quad (10)$$

Assume the following.

$$\forall X0.(((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_finseq\_1 X0)))\Rightarrow((v1\_relat\_1 X0)\wedge((v4\_relat\_1 X0 k5\_numbers)\wedge((v1\_funct\_1 X0)\wedge(v1\_finseq\_1 X0)))))) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_1 X1 X0)\Rightarrow(v5\_relat\_1 X1 X0) \quad (12)$$

**Theorem 1**

$$\forall X0.(((v1\_relat\_1 X0)\wedge((v4\_relat\_1 X0 k5\_numbers)\wedge((v1\_funct\_1 X0)\wedge((v1\_finset\_1 X0)\wedge(v1\_glib\_000 X0))))))\Rightarrow(\forall X1.(m3\_glib\_001 X1 X0)\Rightarrow(\forall X2.(m1\_subset\_1 X2 k5\_numbers)\Rightarrow((X2 \in k1\_relset\_1 k5\_numbers (k12\_glib\_001 X0 X1))\Rightarrow(k1\_funct\_1 (k12\_glib\_001 X0 X1) X2 \in k7\_glib\_000 X0))))))$$