

t79_sheffer2

(TMMwEyUxG8YUN2PSr8D1X5SEdDsCBF1Evhh)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v10_sheffer1 : \iota \Rightarrow o$ be given. Let $v11_sheffer1 : \iota \Rightarrow o$ be given. Let $v12_sheffer1 : \iota \Rightarrow o$ be given. Let $l1_sheffer1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k5_sheffer1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0. (&(\neg v2_struct_0 X0) \wedge ((v10_sheffer1 X0) \wedge ((v11_sheffer1 \\ &X0) \wedge ((v12_sheffer1 X0) \wedge (l1_sheffer1 X0)))))) \Rightarrow (\forall X1. (m1_subset_1 \\ &X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. (m1_subset_1 X2 (u1_struct_0 \\ &X0)) \Rightarrow (X2 = k5_sheffer1 X0 (k5_sheffer1 X0 X2 X2) (k5_sheffer1 X0 \\ &X1 (k5_sheffer1 X0 X1 X1)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0. (&(\neg v2_struct_0 X0) \wedge ((v10_sheffer1 X0) \wedge ((v11_sheffer1 \\ &X0) \wedge ((v12_sheffer1 X0) \wedge (l1_sheffer1 X0)))))) \Rightarrow (\forall X1. (m1_subset_1 \\ &X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. (m1_subset_1 X2 (u1_struct_0 \\ &X0)) \Rightarrow (k5_sheffer1 X0 X2 (k5_sheffer1 X0 (k5_sheffer1 X0 X1 X1) X1) = \\ &k5_sheffer1 X0 X2 X2))) \end{aligned} \tag{2}$$

Theorem 1

$$\begin{aligned} \forall X0. (&(\neg v2_struct_0 X0) \wedge ((v10_sheffer1 X0) \wedge ((v11_sheffer1 \\ &X0) \wedge ((v12_sheffer1 X0) \wedge (l1_sheffer1 X0)))))) \Rightarrow (\forall X1. (m1_subset_1 \\ &X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. (m1_subset_1 X2 (u1_struct_0 \\ &X0)) \Rightarrow (\forall X3. (m1_subset_1 X3 (u1_struct_0 X0)) \Rightarrow (k5_sheffer1 \\ &X0 (k5_sheffer1 X0 X3 (k5_sheffer1 X0 (k5_sheffer1 X0 X2 X2) X2)) \\ &(k5_sheffer1 X0 X1 (k5_sheffer1 X0 X1 X1)) = X3)))) \end{aligned}$$