

t7\_afinsq\_1 (TMdm-  
ngyv6fRKxM8vARQeJf7aWkT1goMNBot)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v5\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finset\_1 : \iota \Rightarrow o$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k2\_afinsq\_1 : \iota \Rightarrow \iota$  be given. Let  $k4\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k2\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $v6\_membered : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1\_subset\_1 X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1\_relat\_1 X2) \wedge (v1\_funct\_1 X2)) \Rightarrow ((k4\_tarski X0 X1 \in X2) \Leftrightarrow ((X0 \in k9\_xtuple\_0 X2) \wedge (X1 = k1\_funct\_1 X2 X0))) \quad (2)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v5\_ordinal1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finset\_1 X0)))) \Rightarrow (k2\_afinsq\_1 X0 = k9\_xtuple\_0 X0) \quad (3)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v5\_ordinal1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finset\_1 X0)))) \Rightarrow (m1\_subset\_1 (k2\_afinsq\_1 X0) (k1\_zfmisc\_1 k5\_numbers)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. k4\_tarski X0 X1 = k2\_tarski (k2\_tarski X0 X1) (k1\_tarski X0) \quad (5)$$

Assume the following.

$$\forall X0. (v1\_relat\_1 X0) \Leftrightarrow (\forall X1. \neg (X1 \in X0) \wedge (\forall X2. \forall X3. X1 \neq k4\_tarski X2 X3)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.k2\_tarSKI X0 X1 = k2\_tarSKI X1 X0 \quad (7)$$

Assume the following.

$$\forall X0.(v6\_membered X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 X0) \Rightarrow (v7\_ordinal1 X1)) \quad (8)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 (k1\_zfmisc\_1 k5\_numbers)) \Rightarrow (v6\_membered X0) \quad (9)$$

**Theorem 1**

$$\forall X0.\forall X1.((v1\_relat\_1 X1) \wedge ((v5\_ordinal1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finset\_1 X1)))) \Rightarrow (\neg(X0 \in X1) \wedge (\forall X2.(v7\_ordinal1 X2) \Rightarrow (\neg(X2 \in k2\_afinsq\_1 X1) \wedge (X0 = k4\_tarSKI X2 (k1\_funct\_1 X1 X2)))))$$