

## t7\_bcialg\_5

(TMQYR4QpbZ5CLFFMVszgnj7GzK9sDHtnrAz)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v3\_bcialg\_1 : \iota \Rightarrow o$  be given. Let  $v4\_bcialg\_1 : \iota \Rightarrow o$  be given. Let  $v5\_bcialg\_1 : \iota \Rightarrow o$  be given. Let  $v7\_bcialg\_1 : \iota \Rightarrow o$  be given. Let  $l2\_bcialg\_1 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_bcialg\_5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k1\_bcialg\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k1\_bcialg\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $np\_0 : \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k2\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v3\_bcialg\_1 X0) \wedge ((v4\_bcialg\_1 \\ & X0) \wedge ((v5\_bcialg\_1 X0) \wedge ((v7\_bcialg\_1 X0) \wedge (l2\_bcialg\_1 X0)))))) \Rightarrow \\ & (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 X0)) \Rightarrow (\forall X2.(m1\_subset\_1 \\ & X2 (u1\_struct\_0 X0)) \Rightarrow (k1\_bcialg\_2 X0 X1 X2 np\_1 = k1\_bcialg\_1 X0 \\ & X1 X2))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v3\_bcialg\_1 X0) \wedge ((v4\_bcialg\_1 \\ & X0) \wedge ((v5\_bcialg\_1 X0) \wedge ((v7\_bcialg\_1 X0) \wedge (l2\_bcialg\_1 X0)))))) \Rightarrow \\ & (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 X0)) \Rightarrow (\forall X2.(m1\_subset\_1 \\ & X2 (u1\_struct\_0 X0)) \Rightarrow (k1\_bcialg\_2 X0 X1 X2 k6\_numbers = X1))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$(m2\_subset\_1\ np\_0\ k1\_numbers\ k5\_numbers) \wedge ((m1\_subset\_1\ np\_0\ k5\_numbers) \wedge (m1\_subset\_1\ np\_0\ k1\_numbers)) \quad (5)$$

Assume the following.

$$v1\_xboole\_0\ np\_0 \quad (6)$$

Assume the following.

$$k2\_xcmplx\_0\ np\_0\ np\_1 = np\_1 \quad (7)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (8)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1\ X0\ k5\_numbers) \wedge (v7\_ordinal1\ X1)) \Rightarrow (k2\_nat\_1\ X0\ X1 = k2\_xcmplx\_0\ X0\ X1) \quad (10)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.(((\neg v2\_struct\_0 \\ & X0) \wedge ((v3\_bcialg\_1\ X0) \wedge ((v4\_bcialg\_1\ X0) \wedge ((v5\_bcialg\_1\ X0) \wedge \\ & ((v7\_bcialg\_1\ X0) \wedge (l2\_bcialg\_1\ X0)))))) \wedge ((m1\_subset\_1\ X1\ (u1\_struct\_0 \\ & X0)) \wedge ((m1\_subset\_1\ X2\ (u1\_struct\_0\ X0)) \wedge (m1\_subset\_1\ X3\ k5\_numbers)))) \Rightarrow \\ & (m1\_subset\_1\ (k1\_bcialg\_2\ X0\ X1\ X2\ X3)\ (u1\_struct\_0\ X0)) \end{aligned} \quad (11)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2\_struct\_0\ X0) \wedge ((v3\_bcialg\_1\ X0) \wedge ((v4\_bcialg\_1 \\ & X0) \wedge ((v5\_bcialg\_1\ X0) \wedge ((v7\_bcialg\_1\ X0) \wedge (l2\_bcialg\_1\ X0)))))) \Rightarrow \\ & (\forall X1.(m1\_subset\_1\ X1\ (u1\_struct\_0\ X0)) \Rightarrow (\forall X2.(m1\_subset\_1 \\ & X2\ (u1\_struct\_0\ X0)) \Rightarrow (\forall X3.(m2\_subset\_1\ X3\ k1\_numbers\ k5\_numbers) \Rightarrow \\ & (\forall X4.(m2\_subset\_1\ X4\ k1\_numbers\ k5\_numbers) \Rightarrow (k1\_bcialg\_5 \\ & X0\ X1\ X2\ X3\ X4 = k1\_bcialg\_2\ X0\ (k1\_bcialg\_2\ X0\ X1\ (k1\_bcialg\_1\ X0\ X1 \\ & X2)\ (k2\_nat\_1\ X3\ np\_1))\ (k1\_bcialg\_1\ X0\ X2\ X1)\ X4)))))) \end{aligned} \quad (12)$$

Assume the following.

$$\forall X0.(m1\_subset\_1\ X0\ k4\_ordinal1) \Rightarrow (v7\_ordinal1\ X0) \quad (13)$$

### Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2\_struct\_0\ X0) \wedge ((v3\_bcialg\_1\ X0) \wedge ((v4\_bcialg\_1 \\ & X0) \wedge ((v5\_bcialg\_1\ X0) \wedge ((v7\_bcialg\_1\ X0) \wedge (l2\_bcialg\_1\ X0)))))) \Rightarrow \\ & (\forall X1.(m1\_subset\_1\ X1\ (u1\_struct\_0\ X0)) \Rightarrow (\forall X2.(m1\_subset\_1 \\ & X2\ (u1\_struct\_0\ X0)) \Rightarrow (k1\_bcialg\_5\ X0\ X1\ X2\ k6\_numbers\ k6\_numbers = \\ & k1\_bcialg\_1\ X0\ X1\ (k1\_bcialg\_1\ X0\ X1\ X2)))) \end{aligned}$$