

t7_boolmark
(TMJnjmDBjEos42Gk6jze7HNx48D4gjnQ7fz)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(\neg v1_xboole_0 X1) \Rightarrow (\\ & \quad \forall X2.(m2_finseq_1 X2 X1) \Rightarrow (\forall X3.(m2_finseq_1 X3 X1) \Rightarrow \\ & ((X0 \in k4_finseq_1 X2) \Rightarrow (k7_partfun1 X1 (k8_finseq_1 X1 X2 X3) X0 = \\ & \quad k7_partfun1 X1 X2 X0)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.(v7_ordinal1 X1) \Rightarrow ((X1 \in k1_relset_1 k5_numbers X0) \Leftrightarrow \\ & ((r1_xxreal_0 np_1 X1) \wedge (r1_xxreal_0 X1 (k3_finseq_1 X0)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \tag{3}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (k4_finseq_1 X0 = k9_xtuple_0 X0) \end{aligned} \tag{5}$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Rightarrow((v1_funct_1 X1)\wedge(v1_finseq_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge(v1_funct_1 X1)\wedge(v1_finseq_1 X1)) \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (9)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow((v1_relat_1 X0)\wedge((v4_relat_1 X0 k5_numbers)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))) \quad (10)$$

Theorem 1

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.(m2_finseq_1 X1 X0)\Rightarrow(\forall X2.(m2_finseq_1 X2 X0)\Rightarrow(\forall X3.(m1_subset_1 X3 k5_numbers)\Rightarrow(((r1_xxreal_0 np_1 X3)\wedge(r1_xxreal_0 X3 (k3_finseq_1 X1)))\Rightarrow(k7_partfun1 X0 (k8_finseq_1 X0 X1 X2) X3 = k7_partfun1 X0 X1 X3))))))$$