

t7\_calcul\_1 (TMS-  
FtXhKXu7cVXAAPKp7WXvzsSPyxB9ZFSa)

October 27, 2020

Let  $m1\_qc\_lang1 : \iota \Rightarrow o$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_cqc\_lang : \iota \Rightarrow \iota$  be given. Let  $k15\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k8\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $v2\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $m1\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ (\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \Rightarrow (X0 = k5\_relat\_1 (k7\_finseq\_1 X0 X1) (k4\_finseq\_1 X0))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v2\_finseq\_1 X0))) \Rightarrow \\ (((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (k15\_finseq\_1 X0 = X0)) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m2\_finseq\_1 X1 X0) \Leftrightarrow (m1\_finseq\_1 X1 X0) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((m1\_finseq\_1 X1 X0) \wedge (m1\_finseq\_1 X2 X0)) \Rightarrow (k8\_finseq\_1 X0 X1 X2 = k7\_finseq\_1 X1 X2) \quad (4)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (k4\_finseq\_1 X0 = k9\_xtuple\_0 X0) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((v1\_funct\_1 X2) \wedge \\ & (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))) \Rightarrow (k2\_partfun1 \\ & X0 X1 X2 X3 = k5\_relat\_1 X2 X3) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow ( \\ & k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m2\_finseq\_1 X1 X0) \Rightarrow ((v1\_funct\_1 X1) \wedge ( \\ & (v1\_finseq\_1 X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers \\ & X0)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1\_finseq\_1 X1 X0) \Rightarrow ((v1\_relat\_1 X1) \wedge ( \\ & (v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((m1\_finseq\_1 X1 X0) \wedge (m1\_finseq\_1 \\ & X2 X0)) \Rightarrow (m2\_finseq\_1 (k8\_finseq\_1 X0 X1 X2) X0) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ & ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v2\_finseq\_1 X0))) \end{aligned} \quad (11)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ & ((v1\_relat\_1 X0) \wedge ((v4\_relat\_1 X0 k5\_numbers) \wedge ((v1\_funct\_1 X0) \wedge \\ & (v1\_finseq\_1 X0)))) \end{aligned} \quad (12)$$

**Theorem 1**

$$\begin{aligned} & \forall X0. (m1\_qc\_lang1 X0) \Rightarrow (\forall X1. (m2\_finseq\_1 X1 (k3\_cqc\_lang \\ & X0)) \Rightarrow (\forall X2. (m2\_finseq\_1 X2 (k3\_cqc\_lang X0)) \Rightarrow (k15\_finseq\_1 \\ & (k2\_partfun1 k5\_numbers (k3\_cqc\_lang X0) (k8\_finseq\_1 (k3\_cqc\_lang \\ & X0) X1 X2) (k1\_relset\_1 k5\_numbers X1)) = k2\_partfun1 k5\_numbers \\ & (k3\_cqc\_lang X0) (k8\_finseq\_1 (k3\_cqc\_lang X0) X1 X2) (k1\_relset\_1 \\ & k5\_numbers X1)))) \end{aligned}$$