

t7_card_3

(TMaM6pWS4LDdYmRVRoR2gDwcsWJc1msHctM)

October 27, 2020

Let $k1_xboole_0 : \iota$ be given. Let $k3_card_3 : \iota \Rightarrow \iota$ be given. Let $k7_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_tarski : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (X1 \in X0) \Rightarrow (k1_funct_1 (k2_funcop_1 X0 X2) X1 = X2) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (r1_tarski X0 (k3_tarski X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. r1_tarski (k3_card_3 (k7_funcop_1 X0 X1)) X1 \quad (3)$$

Assume the following.

$$\forall X0. (v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (k9_xtuple_0 (k2_funcop_1 X0 X1) = X0) \wedge (r1_tarski (k10_xtuple_0 (k2_funcop_1 X0 X1)) (k1_tarski X1)) \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. k7_funcop_1 X0 X1 = k2_funcop_1 X0 X1 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.v1_relat_1 (k2_zfmisc_1 X0 X1) \quad (8)$$

Assume the following.

$$\forall X0.\exists X1.m1_subset_1 X1 X0 \quad (9)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.(v1_funct_1 (k7_funcop_1 X0 X1)) \wedge ((v1_funct_2 \\ (k7_funcop_1 X0 X1) X0 (k1_tarski X1)) \wedge (m1_subset_1 (k7_funcop_1 \\ X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 (k1_tarski X1)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (k3_card_3 X0 = k3_tarski (k10_xtuple_0 X0)) \quad (11)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.(X1 = \\ k10_xtuple_0 X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (\exists X3.(X3 \in k9_xtuple_0 \\ X0) \wedge (X2 = k1_funct_1 X0 X3)))) \end{aligned} \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.k2_funcop_1 X0 X1 = k2_zfmisc_1 X0 (k1_tarski X1) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.(X0 = X1) \Leftrightarrow ((r1_tarski X0 X1) \wedge (r1_tarski X1 X0)) \quad (14)$$

Theorem 1

$$\forall X0.\forall X1.(X0 \neq k1_xboole_0) \Rightarrow (k3_card_3 (k7_funcop_1 X0 X1) = X1)$$