

t7_cfuncdom

(TMPsb6Sjt9D93Us79mFDCCGpuN9SqmdQMBjM)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m2_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_numbers : \iota$ be given. Let $k9_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funcsdom : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_cfuncdom : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_complex1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_binop_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m2_funct_2 X1 X0 k2_numbers \\ & (k9_funct_2 X0 k2_numbers)) \Rightarrow (\forall X2. (m2_funct_2 X2 X0 k2_numbers \\ & (k9_funct_2 X0 k2_numbers)) \Rightarrow (\forall X3. (m2_funct_2 X3 X0 k2_numbers \\ & (k9_funct_2 X0 k2_numbers)) \Rightarrow ((r2_funct_2 X0 k2_numbers X1 (k1_funcsdom \\ & X0 k2_numbers (k2_cfuncdom X0) X2 X3)) \Leftrightarrow (\forall X4. (m1_subset_1 \\ & X4 X0) \Rightarrow (k3_funct_2 X0 k2_numbers X1 X4 = k9_complex1 (k3_funct_2 \\ & X0 k2_numbers X2 X4) (k3_funct_2 X0 k2_numbers X3 X4))))))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. (((v1_funct_1 X2) \wedge \\ & ((v1_funct_2 X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 X1)))))) \wedge ((v1_funct_1 X3) \wedge ((v1_funct_2 X3 X0 X1) \wedge (m1_subset_1 \\ & X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \Rightarrow (r2_funct_2 X0 X1 X2 X2) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((\neg v1_xboole_0 X1) \wedge (m1_funct_2 \\ & X2 X0 X1)) \Rightarrow (\forall X3. (m2_funct_2 X3 X0 X1 X2) \Leftrightarrow (m1_subset_1 X3 \\ & X2)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (\neg v1_xboole_0 X1) \Rightarrow (k9_funct_2 X0 X1 = k1_funct_2 \\ & X0 X1) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.\forall X4.((\neg v1_xboole_0 \\ & X1)\wedge(((v1_funct_1 X2)\wedge((v1_funct_2 X2 (k2_zfmisc_1 (k9_funct_2 \\ & X0 X1) (k9_funct_2 X0 X1)) (k9_funct_2 X0 X1))\wedge(m1_subset_1 X2 (\\ & k1_zfmisc_1 (k2_zfmisc_1 (k2_zfmisc_1 (k9_funct_2 X0 X1) (k9_funct_2 \\ & X0 X1)) (k9_funct_2 X0 X1))))))\wedge((m1_subset_1 X3 (k9_funct_2 X0 \\ & X1))\wedge(m1_subset_1 X4 (k9_funct_2 X0 X1))))))\Rightarrow(k1_funcsdm X0 X1 \\ & X2 X3 X4 = k1_binop_1 X2 X3 X4) \end{aligned} \quad (5)$$

Assume the following.

$$\neg v1_xboole_0 k2_numbers \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X1)\wedge(m1_funct_2 \\ & X2 X0 X1))\Rightarrow(\forall X3.(m2_funct_2 X3 X0 X1 X2)\Rightarrow((v1_funct_1 X3)\wedge \\ & ((v1_funct_2 X3 X0 X1)\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 X1)))))) \end{aligned} \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(\neg v1_xboole_0 X1)\Rightarrow(m1_funct_2 (k9_funct_2 X0 X1) X0 X1) \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0)\wedge \\ & (((v1_funct_1 X2)\wedge((v1_funct_2 X2 X0 X1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))))))\wedge(m1_subset_1 X3 X0))\Rightarrow(m1_subset_1 (\\ & k3_funct_2 X0 X1 X2 X3) X1) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0)\Rightarrow((v1_funct_1 (k2_cfundom X0))\wedge \\ & ((v1_funct_2 (k2_cfundom X0) (k2_zfmisc_1 (k9_funct_2 X0 k2_numbers) \\ & (k9_funct_2 X0 k2_numbers)) (k9_funct_2 X0 k2_numbers))\wedge(m1_subset_1 \\ & (k2_cfundom X0) (k1_zfmisc_1 (k2_zfmisc_1 (k2_zfmisc_1 (k9_funct_2 \\ & X0 k2_numbers) (k9_funct_2 X0 k2_numbers)) (k9_funct_2 X0 k2_numbers)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.\forall X4.((\neg v1_xboole_0 \\ & X1)\wedge(((v1_funct_1 X2)\wedge((v1_funct_2 X2 (k2_zfmisc_1 (k9_funct_2 \\ & X0 X1) (k9_funct_2 X0 X1)) (k9_funct_2 X0 X1))\wedge(m1_subset_1 X2 (\\ & k1_zfmisc_1 (k2_zfmisc_1 (k2_zfmisc_1 (k9_funct_2 X0 X1) (k9_funct_2 \\ & X0 X1)) (k9_funct_2 X0 X1))))))\wedge((m1_subset_1 X3 (k9_funct_2 X0 \\ & X1))\wedge(m1_subset_1 X4 (k9_funct_2 X0 X1))))))\Rightarrow(m2_funct_2 (k1_funcsdm \\ & X0 X1 X2 X3 X4) X0 X1 (k9_funct_2 X0 X1)) \end{aligned} \quad (11)$$

Assume the following.

$$\forall X0. \forall X1. ((m1_subset_1 X0 k2_numbers) \wedge (m1_subset_1 X1 k2_numbers)) \Rightarrow (k9_complex1 X0 X1 = k9_complex1 X1 X0) \quad (12)$$

Theorem 1

$$\begin{aligned} \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m2_funct_2 X1 X0 k2_numbers \\ (k9_funct_2 X0 k2_numbers)) \Rightarrow (\forall X2. (m2_funct_2 X2 X0 k2_numbers \\ (k9_funct_2 X0 k2_numbers)) \Rightarrow (r2_funct_2 X0 k2_numbers (k1_funcsdom \\ X0 k2_numbers (k2_cfunctor X0) X1 X2) (k1_funcsdom X0 k2_numbers \\ (k2_cfunctor X0) X2 X1)))) \end{aligned}$$