

t7_circled1 (TMHX-
mAi65hR47oABJyj5tzK1EtFbLAbXWkN)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rlvect_1 : \iota \Rightarrow o$ be given. Let $v6_rlvect_1 : \iota \Rightarrow o$ be given. Let $v7_rlvect_1 : \iota \Rightarrow o$ be given. Let $v8_rlvect_1 : \iota \Rightarrow o$ be given. Let $l1_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rltopsp1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k18_complex1 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_rlvect_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow (k9_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow (m1_subset_1 (k9_subset_1 X0 X1 X2) (k1_zfmisc_1 X0)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k3_xboole_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (X3 \in X1))) \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge (l1_rlvect_1 X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow ((v3_rltopsp1 \\ & X1 X0) \Leftrightarrow (\forall X2. (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (\forall X3. \\ & (m1_subset_1 X3 k1_numbers) \Rightarrow (((r1_xxreal_0 (k18_complex1 X3) \\ & np_1) \wedge (X2 \in X1)) \Rightarrow (k1_rlvect_1 X0 X2 X3 \in X1)))))) \quad (4) \end{aligned}$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (v13_algstr_0 X0) \wedge (v2_rlvect_1 \\ & X0) \wedge (v3_rlvect_1 X0) \wedge (v4_rlvect_1 X0) \wedge (v5_rlvect_1 X0) \wedge \\ & ((v6_rlvect_1 X0) \wedge (v7_rlvect_1 X0) \wedge (v8_rlvect_1 X0) \wedge (l1_rlvect_1 \\ & X0)))))) \Rightarrow (\forall X1.((v3_rltopsp1 X1 X0) \wedge (m1_subset_1 \\ & X1 (k1_zfmisc_1 (u1_struct_0 X0)))) \Rightarrow (\forall X2.((v3_rltopsp1 \\ & X2 X0) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X0)))) \Rightarrow (v3_rltopsp1 \\ & (k9_subset_1 (u1_struct_0 X0) X1 X2) X0))) \end{aligned}$$