

t7\_complex1  
(TMYVskNpUJtzDdYofU8CRFSvuxgnRSxYDjr)

October 27, 2020

Let  $k3\_complex1 : \iota \Rightarrow \iota$  be given. Let  $k7\_complex1 : \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k4\_complex1 : \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_arytm\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (\forall X1.(m1\_subset\_1 \\ X1 k1\_numbers) \Rightarrow ((k3\_complex1 (k5\_arytm\_0 X0 X1) = X0) \wedge (k4\_complex1 \\ (k5\_arytm\_0 X0 X1) = X1))) \end{aligned} \tag{1}$$

Assume the following.

$$m1\_subset\_1 np\_1 k1\_numbers \tag{2}$$

Assume the following.

$$m1\_subset\_1 k6\_numbers k1\_numbers \tag{3}$$

Assume the following.

$$k7\_complex1 = k5\_arytm\_0 k6\_numbers np\_1 \tag{4}$$

**Theorem 1**

$$(k3\_complex1 k7\_complex1 = k6\_numbers) \wedge (k4\_complex1 k7\_complex1 = np\_1)$$