

t7_diff_2

(TMRrtPq2Pje4Am1xMwd1aty3E8o5ySYrifE)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_diff_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k26_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k24_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (\forall X1.(m1_subset_1 \\
 & \quad X1 k1_numbers) \Rightarrow (\forall X2.(m1_subset_1 X2 k1_numbers) \Rightarrow (\forall X3. \\
 & ((v1_funct_1 X3) \wedge ((v1_funct_2 X3 k1_numbers k1_numbers) \wedge (m1_subset_1 \\
 & \quad X3 (k1_zfmisc_1 (k2_zfmisc_1 k1_numbers k1_numbers)))))) \Rightarrow (\forall X4. \\
 & ((v1_funct_1 X4) \wedge ((v1_funct_2 X4 k1_numbers k1_numbers) \wedge (m1_subset_1 \\
 & \quad X4 (k1_zfmisc_1 (k2_zfmisc_1 k1_numbers k1_numbers)))))) \Rightarrow (k10_diff_1 \\
 & (k3_valued_1 k1_numbers k1_numbers k1_numbers X3 X4) X0 X1 X2 = k7_real_1 \\
 & \quad (k10_diff_1 X3 X0 X1 X2) (k10_diff_1 X4 X0 X1 X2))))))
 \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
 & \forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (\forall X1.(m1_subset_1 \\
 & \quad X1 k1_numbers) \Rightarrow (\forall X2.(m1_subset_1 X2 k1_numbers) \Rightarrow (\forall X3. \\
 & \quad (m1_subset_1 X3 k1_numbers) \Rightarrow (\forall X4.((v1_funct_1 X4) \wedge ((\\
 & v1_funct_2 X4 k1_numbers k1_numbers) \wedge (m1_subset_1 X4 (k1_zfmisc_1 \\
 & \quad (k2_zfmisc_1 k1_numbers k1_numbers)))))) \Rightarrow (k10_diff_1 (k26_valued_1 \\
 & k1_numbers k1_numbers X4 X0) X1 X2 X3 = k8_real_1 X0 (k10_diff_1 X4 \\
 & \quad X1 X2 X3))))))
 \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned}
 & \forall X0.\forall X1.\forall X2.\forall X3.((v3_membered X1) \wedge \\
 & (((v1_funct_1 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\
 & X0 X1)))) \wedge (v1_xreal_0 X3))) \Rightarrow (k26_valued_1 X0 X1 X2 X3 = k24_valued_1 \\
 & \quad X2 X3)
 \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.(((\neg v1_xboole_0 \\ & X1)\wedge(v3_membered\ X1))\wedge(((v1_funct_1\ X2)\wedge((v1_funct_2\ X2\ X0\ X1)\wedge \\ & (m1_subset_1\ X2\ (k1_zfmisc_1\ (k2_zfmisc_1\ X0\ X1))))))\wedge(v1_xreal_0 \\ & X3)))\Rightarrow((v1_funct_1\ (k24_valued_1\ X2\ X3))\wedge(v1_partfun1\ (k24_valued_1 \\ & X2\ X3)\ X0)) \end{aligned} \tag{4}$$

Assume the following.

$$v3_membered\ k1_numbers \tag{5}$$

Assume the following.

$$\neg v1_xboole_0\ k1_numbers \tag{6}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((v3_membered\ X1)\wedge \\ & (((v1_funct_1\ X2)\wedge(m1_subset_1\ X2\ (k1_zfmisc_1\ (k2_zfmisc_1 \\ & X0\ X1))))\wedge(v1_xreal_0\ X3)))\Rightarrow((v1_funct_1\ (k26_valued_1\ X0\ X1 \\ & X2\ X3))\wedge(m1_subset_1\ (k26_valued_1\ X0\ X1\ X2\ X3)\ (k1_zfmisc_1\ (k2_zfmisc_1 \\ & X0\ k1_numbers)))) \end{aligned} \tag{7}$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k1_numbers)\Rightarrow(v1_xreal_0\ X0) \tag{8}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1_subset_1\ X2\ (k1_zfmisc_1 \\ & (k2_zfmisc_1\ X0\ X1)))\Rightarrow((v1_partfun1\ X2\ X0)\Rightarrow(v1_funct_2\ X2\ X0\ X1)) \end{aligned} \tag{9}$$

Theorem 1

$$\begin{aligned} & \forall X0.(m1_subset_1\ X0\ k1_numbers)\Rightarrow(\forall X1.(m1_subset_1 \\ & X1\ k1_numbers)\Rightarrow(\forall X2.(m1_subset_1\ X2\ k1_numbers)\Rightarrow(\forall X3. \\ & (m1_subset_1\ X3\ k1_numbers)\Rightarrow(\forall X4.(m1_subset_1\ X4\ k1_numbers)\Rightarrow \\ & (\forall X5.((v1_funct_1\ X5)\wedge((v1_funct_2\ X5\ k1_numbers\ k1_numbers)\wedge \\ & (m1_subset_1\ X5\ (k1_zfmisc_1\ (k2_zfmisc_1\ k1_numbers\ k1_numbers))))))\Rightarrow \\ & (\forall X6.((v1_funct_1\ X6)\wedge((v1_funct_2\ X6\ k1_numbers\ k1_numbers)\wedge \\ & (m1_subset_1\ X6\ (k1_zfmisc_1\ (k2_zfmisc_1\ k1_numbers\ k1_numbers))))))\Rightarrow \\ & (k10_diff_1\ (k3_valued_1\ k1_numbers\ k1_numbers\ k1_numbers\ (k26_valued_1 \\ & k1_numbers\ k1_numbers\ X5\ X0)\ (k26_valued_1\ k1_numbers\ k1_numbers \\ & X6\ X1))\ X2\ X3\ X4 = k7_real_1\ (k8_real_1\ X0\ (k10_diff_1\ X5\ X2\ X3\ X4) \\ & (k8_real_1\ X1\ (k10_diff_1\ X6\ X2\ X3\ X4)))))) \end{aligned}$$