

t7_euclid_4 (TM- RXmZwxapBBK3R2GFouiBkEpLb6Z14wJHr)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $m2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_euclid : \iota \Rightarrow \iota$ be given. Let $k9_euclid : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_euclid : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k10_rvsum_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_binop_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_rvsum_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_finseq_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k24_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & ((v1_relat_1 X2) \wedge ((v1_funct_1 X2) \wedge ((v3_valued_0 X2) \wedge (v1_finseq_1 \\ & X2)))) \Rightarrow (k10_rvsum_1 X2 (k9_binop_2 X0 X1) = k4_rvsum_1 (k10_rvsum_1 \\ & X2 X0) (k10_rvsum_1 X2 X1)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_2 X1 X0) \Rightarrow (\forall X2.(m2_finseq_2 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \tag{2}$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((v7_ordinal1 X0) \wedge ((m1_subset_1 \\ & X1 (k1_euclid X0)) \wedge (v1_xreal_0 X2))) \Rightarrow (k9_euclid X0 X1 X2 = k24_valued_1 \\ & X1 X2) \end{aligned} \tag{4}$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow(k9_binop_2 X0 X1 = k2_xcmplx_0 X0 X1) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers)\wedge(v1_xreal_0 X1))\Rightarrow(k7_real_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v7_ordinal1 X0)\wedge((m1_subset_1 X1 (k1_euclid X0))\wedge(m1_subset_1 X2 (k1_euclid X0))))\Rightarrow(k7_euclid X0 X1 X2 = k1_valued_1 X1 X2) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge((v3_valued_0 X0)\wedge(v1_finseq_1 X0))))\wedge((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge((v3_valued_0 X1)\wedge(v1_finseq_1 X1)))))\Rightarrow(k4_rvsum_1 X0 X1 = k1_valued_1 X0 X1) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge((v3_valued_0 X0)\wedge(v1_finseq_1 X0))))\wedge(v1_xreal_0 X1))\Rightarrow(k10_rvsum_1 X0 X1 = k24_valued_1 X0 X1) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow(v1_xreal_0 (k2_xcmplx_0 X0 X1)) \quad (10)$$

Assume the following.

$$v3_membered k1_numbers \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_2 X1 X0)\Rightarrow(\forall X2.(m2_finseq_2 X2 X0 X1)\Rightarrow(m2_finseq_1 X2 X0)) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Rightarrow((v1_funct_1 X1)\wedge((v1_finseq_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge(v1_funct_1 X1)\wedge(v1_finseq_1 X1)) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v7_ordinal1 X0)\wedge((m1_subset_1 X1 (k1_euclid X0))\wedge(v1_xreal_0 X2)))\Rightarrow(m2_finseq_2 (k9_euclid X0 X1 X2) k1_numbers (k1_euclid X0)) \quad (15)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(m1_finseq_2 (k1_euclid X0) k1_numbers) \quad (16)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.(v3_membered X1)\Rightarrow(\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v3_valued_0 X2)) \quad (18)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(\forall X1.(m1_subset_1 X1 k1_numbers)\Rightarrow(\forall X2.(v7_ordinal1 X2)\Rightarrow(\forall X3.(m2_finseq_2 X3 k1_numbers (k1_euclid X2))\Rightarrow(k9_euclid X2 X3 (k7_real_1 X0 X1) = k7_euclid X2 (k9_euclid X2 X3 X0) (k9_euclid X2 X3 X1))))))$$