

t7_fin_topo
(TMaQoxv2EfphPysNmpdaump7cN9z2LuxvRh)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_orders_2 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k8_fin_topo : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_fin_topo : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge (l1_orders_2 X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (k8_fin_topo \\ & X0 X1 = ReplSep (toset (\lambda X2 : \iota. m1_subset_1 X2 (u1_struct_0 \\ & X0)))) (\lambda X2 : \iota. r1_tarski (k1_fin_topo X0 X2) X1) (\lambda X2 : \iota. \\ & X2))) \end{aligned} \quad (1)$$

Theorem 1

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge (l1_orders_2 X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. (m1_subset_1 X2 \\ & (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow ((X1 \in k8_fin_topo X0 X2) \Leftrightarrow (r1_tarski \\ & (k1_fin_topo X0 X1) X2)))) \end{aligned}$$