

t7_finseq_1 (TMFWqQPqJzNAVY- Hcn2xcmTaNFAXXEowV8g)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((r1_xreal_0 X0 X1) \Leftrightarrow (r1_tarski (k2_finseq_1 X0) (k2_finseq_1 X1)))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarski X0 X1) \Rightarrow (k3_xboole_0 X0 X1 = X0) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow (k9_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \quad (3)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (m1_subset_1 (k2_finseq_1 X0) (k1_zfmisc_1 k5_numbers)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (5)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((r1_xreal_0 X0 X1) \Rightarrow (k2_finseq_1 X0 = k9_subset_1 k5_numbers (k2_finseq_1 X1) (k2_finseq_1 X0))))$$