

# t7\_finseq\_8 (TMS- GudxJ3VtaULY1dQmDxYJPtAwhTGZJJcF)

October 27, 2020

Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_finseq\_8 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k7\_nat\_d : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $np\_0 : \iota$  be given. Let  $m1\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $k2\_finseq\_8 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k16\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_rfinseq : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow ((r1\_xxreal\_0 X1 X0) \Rightarrow (k7\_nat\_d X1 X0 = k6\_numbers))) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (2)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow ((\neg r1\_xxreal\_0 (k1\_nat\_1 X1 np\_1) X0) \Leftrightarrow (r1\_xxreal\_0 X0 X1))) \quad (3)$$

Assume the following.

$$((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \quad (4)$$

Assume the following.

$$v1\_xboole\_0 np\_0 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0) \Leftrightarrow (m1\_finseq\_1 X1 X0) \quad (6)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (7)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((m1\_finseq\_1 X1 X0)\wedge((v7\_ordinal1 X2)\wedge(v7\_ordinal1 X3)))\Rightarrow(k3\_finseq\_8 X0 X1 X2 X3 = k2\_finseq\_8 X1 X2 X3) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v7\_ordinal1 X0)\wedge(m1\_subset\_1 X1 k5\_numbers))\Rightarrow(k1\_nat\_1 X0 X1 = k2\_xcmplx\_0 X0 X1) \quad (10)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_finseq\_1 X0)))\Rightarrow((v1\_relat\_1 (k16\_finseq\_1 k6\_numbers X0))\wedge((v1\_funct\_1 (k16\_finseq\_1 k6\_numbers X0))\wedge((v1\_xboole\_0 (k16\_finseq\_1 k6\_numbers X0))\wedge(v1\_finseq\_1 (k16\_finseq\_1 k6\_numbers X0)))))) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((v7\_ordinal1 X0)\wedge(v7\_ordinal1 X1))\Rightarrow(v7\_ordinal1 (k2\_xcmplx\_0 X0 X1)) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0)\Rightarrow((v1\_funct\_1 X1)\wedge((v1\_finseq\_1 X1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers X0)))))) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_1 X1 X0)\Rightarrow((v1\_relat\_1 X1)\wedge((v1\_funct\_1 X1)\wedge(v1\_finseq\_1 X1))) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.((v7\_ordinal1 X0)\wedge(v7\_ordinal1 X1))\Rightarrow(m1\_subset\_1 (k7\_nat\_d X0 X1) k5\_numbers) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.((v7\_ordinal1 X0)\wedge((v1\_relat\_1 X1)\wedge((v1\_funct\_1 X1)\wedge(v1\_finseq\_1 X1))))\Rightarrow((v1\_relat\_1 (k1\_rfinseq X0 X1))\wedge((v1\_funct\_1 (k1\_rfinseq X0 X1))\wedge(v1\_finseq\_1 (k1\_rfinseq X0 X1)))) \quad (16)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ (\forall X1.(v7\_ordinal1 X1) \Rightarrow (\forall X2.(v7\_ordinal1 X2) \Rightarrow ( \\ k2\_finseq\_8 X0 X1 X2 = k16\_finseq\_1 (k7\_nat\_d (k1\_nat\_1 X2 np\_1) \\ X1) (k1\_rfinseq (k7\_nat\_d X1 np\_1) X0)))) \end{aligned} \quad (17)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (18)$$

**Theorem 1**

$$\begin{aligned} \forall X0. \forall X1.(m2\_finseq\_1 X1 X0) \Rightarrow (\forall X2.(m1\_subset\_1 \\ X2 k5\_numbers) \Rightarrow (\forall X3.(m1\_subset\_1 X3 k5\_numbers) \Rightarrow ((\neg r1\_xreal\_0 \\ X2 X3) \Rightarrow (k3\_finseq\_8 X0 X1 X2 X3 = k1\_xboole\_0)))) \end{aligned}$$