

t7_funct_1 (TMPWki- AYmm6nHa7x8pZtM9dP1crme2gxZAV)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((\\ v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (((k9_xtuple_0 X0 = k9_xtuple_0 \\ X1) \wedge (\forall X2.(X2 \in k9_xtuple_0 X0) \Rightarrow (k1_funct_1 X0 X2 = k1_funct_1 \\ X1 X2)))) \Rightarrow (X0 = X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.(X1 = \\ k10_xtuple_0 X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (\exists X3.(X3 \in k9_xtuple_0 \\ X0) \wedge (X2 = k1_funct_1 X0 X3)))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1.(X1 = k1_tarski X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (3)$$

Theorem 1

$$\begin{aligned} \forall X0. \forall X1.((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2. \\ ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow (((k9_xtuple_0 X1 = k9_xtuple_0 \\ X2) \wedge ((k10_xtuple_0 X1 = k1_tarski X0) \wedge (k10_xtuple_0 X2 = k1_tarski \\ X0))) \Rightarrow (X1 = X2))) \end{aligned}$$