

t7_jordan7 (TMXKHM- CJdV1Fy3V6nZMJGRLEXwsdpMgoZn8)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v2_compts_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v1_topreal2 : \iota \Rightarrow o$ be given. Let $k1_jordan7 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k18_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $r1_jordan6 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid \\ & \quad np_2)))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid \\ & \quad np_2))) \Rightarrow (((v1_topreal2 X0) \wedge (X1 \in X0)) \Rightarrow (r1_jordan6 X0 X1 X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid \\ & \quad np_2)))) \Rightarrow (m1_subset_1 (k18_pscomp_1 X0) (u1_struct_0 (k15_euclid \\ & \quad np_2))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v1_xboole_0 X0) \wedge ((v2_compts_1 X0 (k15_euclid np_2)) \wedge \\ & \quad (m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))))) \Rightarrow \\ & \quad (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & \quad (\forall X2.(m1_subset_1 X2 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & \quad (((X2 \neq k18_pscomp_1 X0) \Rightarrow (k1_jordan7 X0 X1 X2 = ReplSep (toset (\lambda X3 : \\ & \quad \iota.m1_subset_1 X3 (u1_struct_0 (k15_euclid np_2)))))) (\lambda X3 : \\ & \quad \iota.(r1_jordan6 X0 X1 X3) \wedge (r1_jordan6 X0 X3 X2)) (\lambda X3 : \iota.X3)))) \wedge \\ & \quad ((X2 = k18_pscomp_1 X0) \Rightarrow (k1_jordan7 X0 X1 X2 = ReplSep (toset (\lambda X3 : \\ & \quad \iota.m1_subset_1 X3 (u1_struct_0 (k15_euclid np_2)))))) (\lambda X3 : \\ & \quad \iota.(r1_jordan6 X0 X1 X3) \vee ((X1 \in X0) \wedge (X3 = k18_pscomp_1 X0))) (\lambda X3 : \\ & \quad \iota.X3)))))) \end{aligned} \quad (3)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v1_xboole_0 X0) \wedge ((v2_compts_1 X0 (k15_euclid np_2)) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))))) \Rightarrow \\ & (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (((X1 \in X0) \wedge (v1_topreal2 X0)) \Rightarrow (X1 \in k1_jordan7 X0 X1 (k18_pscomp_1 \\ & X0)))) \end{aligned}$$