

t7_lfuzzy_0

(TMcr1vgiSRxcN8wvQyGsw4CWt2YwQpiNRP6)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v1_lfuzzy_0 : \iota \Rightarrow o$ be given. Let $l1_orders_2 : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $r1_xreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v2_yellow_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_orders_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Let $r2_lattice3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.((\neg v2_struct_0 X0) \wedge ((v1_lfuzzy_0 X0) \wedge (l1_orders_2 \\ X0))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. \\ (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow ((r1_xreal_0 X1 X2) \Leftrightarrow (r1_orders_2 \\ X0 X1 X2)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1.(m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \tag{2}$$

Assume the following.

$$\forall X0. \forall X1.(X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \tag{3}$$

Assume the following.

$$\forall X0.((\neg v2_struct_0 X0) \wedge (l1_struct_0 X0)) \Rightarrow (\neg v1_xboole_0 (u1_struct_0 X0)) \tag{4}$$

Assume the following.

$$\forall X0.(l1_orders_2 X0) \Rightarrow (l1_struct_0 X0) \tag{5}$$

Assume the following.

$$\begin{aligned} \forall X0.(l1_orders_2 X0) \Rightarrow (\forall X1. \forall X2.(m1_subset_1 \\ X2 (u1_struct_0 X0)) \Rightarrow ((r2_lattice3 X0 X1 X2) \Leftrightarrow (\forall X3.(m1_subset_1 \\ X3 (u1_struct_0 X0)) \Rightarrow ((X3 \in X1) \Rightarrow (r1_orders_2 X0 X3 X2)))))) \end{aligned} \tag{6}$$

Assume the following.

$$\forall X0.(l1_orders_2 X0) \Rightarrow ((v2_yellow_0 X0) \Leftrightarrow (\exists X1.(m1_subset_1 X1 (u1_struct_0 X0)) \wedge (r2_lattice3 X0 (u1_struct_0 X0) X1))) \quad (7)$$

Assume the following.

$$\forall X0.((\neg v2_struct_0 X0) \wedge ((v1_lfuzzy_0 X0) \wedge (l1_orders_2 X0))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (v1_xreal_0 X1)) \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v1_lfuzzy_0 X0) \wedge (l1_orders_2 \\ & X0))) \Rightarrow ((\exists X1.(v1_xreal_0 X1) \wedge ((X1 \in u1_struct_0 X0) \wedge (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow ((X2 \in u1_struct_0 X0) \Rightarrow (r1_xxreal_0 X2 X1)))))) \Leftrightarrow \\ & (v2_yellow_0 X0) \end{aligned}$$