

t7_nat_1 (TMNtqBPp-
MQFK8xXqvbFV m5ERdAJV7MzBJBk)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $r1_xreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (r1_xreal_0 k6_numbers X0) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xreal_0 k6_numbers X0) \wedge (r1_xreal_0 k6_numbers X1) \wedge (k2_xcmplx_0 X0 X1 = k6_numbers))) \Rightarrow (X0 = k6_numbers))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (k2_xcmplx_0 X0 X1 = k2_xcmplx_0 X1 X0) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (4)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_xreal_0 X0) \quad (5)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((k2_xcmplx_0 X0 X1 = k6_numbers) \Rightarrow ((X0 = k6_numbers) \wedge (X1 = k6_numbers))))$$