

t7\_ordinal3

(TMZ8iZNqMd1qEwey75M5CDQU9An2DdEcjWq)

October 27, 2020

Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k2\_ordinal1 : \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $k2\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. k2\_tarski X0 X0 = k1\_tarski X0 \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k2\_ordinal1 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow ((X2 \in X0) \wedge (v3\_ordinal1 X2))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1\_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (3)$$

**Theorem 1**

$$\forall X0. (v3\_ordinal1 X0) \Rightarrow (k2\_ordinal1 (k1\_tarski X0) = k1\_tarski X0)$$