

t7_ranknull
(TMVFxpdmj38KwmijiqRg9QpxqLUfqi4Bw96)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_struct_0 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. v1_relat_1 (k2_zfmisc_1 X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. ((\neg v2_struct_0 X0) \wedge (l1_struct_0 X0)) \Rightarrow (\neg v1_xboole_0 (k2_struct_0 X0)) \quad (2)$$

Assume the following.

$$v1_xboole_0 k1_xboole_0 \quad (3)$$

Assume the following.

$$\forall X0. (l1_struct_0 X0) \Rightarrow (k2_struct_0 X0 = u1_struct_0 X0) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))) \Rightarrow (((X1 \neq k1_xboole_0) \Rightarrow ((v1_funct_2 X2 X0 \\ & X1) \Leftrightarrow (X0 = k1_relset_1 X0 X2))) \wedge ((X1 = k1_xboole_0) \Rightarrow ((v1_funct_2 \\ & X2 X0 X1) \Leftrightarrow (X2 = k1_xboole_0)))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow ((v5_relat_1 X1 X0) \Leftrightarrow (r1_tarski (k10_xtuple_0 X1) X0)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (7)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0))\Rightarrow(v1_relat_1 X1)) \quad (8)$$

Theorem 1

$$\begin{aligned} &\forall X0.((\neg v2_struct_0 X0)\wedge(l1_struct_0 X0))\Rightarrow(\forall X1. \\ &((\neg v2_struct_0 X1)\wedge(l1_struct_0 X1))\Rightarrow(\forall X2.((v1_funct_1 \\ &X2)\wedge((v1_funct_2 X2 (u1_struct_0 X0) (u1_struct_0 X1))\wedge(m1_subset_1 \\ &X2 (k1_zfmisc_1 (k2_zfmisc_1 (u1_struct_0 X0) (u1_struct_0 X1))))))\Rightarrow \\ &((k1_relset_1 (u1_struct_0 X0) X2 = k2_struct_0 X0)\wedge(r1_tarski \\ &(k10_xtuple_0 X2) (k2_struct_0 X1)))))) \end{aligned}$$