

t7_rearran1

(TMapuWqas3CwtriUEftSLF2fLTFNJ6QbykY)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_rearran1 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $v3_xxreal_0 : \iota \Rightarrow o$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $np_0 : \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k5_card_1 : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow ((v1_xboole_0 X0) \vee ((v2_xxreal_0 X1) \vee (v3_xxreal_0 X0)))))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (2)$$

Assume the following.

$$k1_card_1 k1_xboole_0 = k1_xboole_0 \quad (3)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((X1 \in k1_relset_1 k5_numbers X0) \Leftrightarrow ((r1_xxreal_0 np_1 X1) \wedge (r1_xxreal_0 X1 (k3_finseq_1 X0)))))) \quad (4)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (5)$$

Assume the following.

$$\neg v1_xboole_0 \ np_1 \tag{6}$$

Assume the following.

$$(m2_subset_1 \ np_0 \ k1_numbers \ k5_numbers) \wedge ((m1_subset_1 \ np_0 \ k5_numbers) \wedge (m1_subset_1 \ np_0 \ k1_numbers)) \tag{7}$$

Assume the following.

$$v1_xboole_0 \ np_0 \tag{8}$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 \ X1 \ X0) \Leftrightarrow (m1_finseq_1 \ X1 \ X0) \tag{9}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{10}$$

Assume the following.

$$\forall X0. (v1_finset_1 \ X0) \Rightarrow (k5_card_1 \ X0 = k1_card_1 \ X0) \tag{11}$$

Assume the following.

$$\forall X0. ((v1_relat_1 \ X0) \wedge ((v1_funct_1 \ X0) \wedge (v1_finseq_1 \ X0))) \Rightarrow (k4_finseq_1 \ X0 = k9_xtuple_0 \ X0) \tag{12}$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 \ X1) \wedge (v4_relat_1 \ X1 \ X0)) \Rightarrow (k1_relset_1 \ X0 \ X1 = k9_xtuple_0 \ X1) \tag{13}$$

Assume the following.

$$v6_membered \ k4_ordinal1 \tag{14}$$

Assume the following.

$$\forall X0. \forall X1. (m1_finseq_1 \ X1 \ X0) \Rightarrow ((v1_relat_1 \ X1) \wedge ((v1_funct_1 \ X1) \wedge (v1_finseq_1 \ X1))) \tag{15}$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 \ X0) \wedge ((v1_funct_1 \ X0) \wedge (v1_finseq_1 \ X0))) \Rightarrow \\ & ((v1_rearran1 \ X0) \Leftrightarrow (\forall X1. (v7_ordinal1 \ X1) \Rightarrow (((r1_xxreal_0 \ np_1 \ X1) \wedge (r1_xxreal_0 \ X1 \ (k3_finseq_1 \ X0))) \Rightarrow (\forall X2. (v1_finset_1 \ X2) \Rightarrow ((X2 = k1_funct_1 \ X0 \ X1) \Rightarrow (k5_card_1 \ X2 = X1)))))) \end{aligned} \tag{16}$$

Assume the following.

$$\forall X0. ((v1_relat_1 \ X0) \wedge ((v1_funct_1 \ X0) \wedge (v1_finseq_1 \ X0))) \Rightarrow ((v1_relat_1 \ X0) \wedge ((v4_relat_1 \ X0 \ k5_numbers) \wedge ((v1_funct_1 \ X0) \wedge (v1_finseq_1 \ X0)))) \tag{17}$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(v1_finset_1\ X0) \quad (18)$$

Assume the following.

$$\forall X0.(v1_xreal_0\ X0)\Rightarrow(v1_xxreal_0\ X0) \quad (19)$$

Assume the following.

$$\forall X0.((v1_xxreal_0\ X0)\wedge(v2_xxreal_0\ X0))\Rightarrow((\neg v1_xboole_0\ X0)\wedge((v1_xxreal_0\ X0)\wedge(\neg v3_xxreal_0\ X0))) \quad (20)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow((v7_ordinal1\ X0)\wedge(\neg v3_xxreal_0\ X0)) \quad (21)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(v1_xreal_0\ X0) \quad (22)$$

Assume the following.

$$\forall X0.(v6_membered\ X0)\Rightarrow(\forall X1.(m1_subset_1\ X1\ X0)\Rightarrow(v7_ordinal1\ X1)) \quad (23)$$

Theorem 1

$$\forall X0.(m1_subset_1\ X0\ k5_numbers)\Rightarrow(\forall X1.((\neg v1_xboole_0\ X1)\wedge(v1_finset_1\ X1))\Rightarrow(\forall X2.((v1_rearran1\ X2)\wedge(m2_finseq_1\ X2\ (k1_zfmisc_1\ X1)))\Rightarrow(\neg(X0 \in k4_finseq_1\ X2)\wedge(k1_funct_1\ X2\ X0 = k1_xboole_0))))$$