

## t7\_relset\_2

(TMSizkbWA7jVsfgReUN5a3V5SdkHszBnqkH)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k8\_setfam\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k1\_zfmisc\_1 \\ & X0))) \Rightarrow (\forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k1\_zfmisc\_1 \\ & X0))) \Rightarrow ((r1\_tarski X1 X2) \Rightarrow (r1\_tarski (k8\_setfam\_1 X0 X2) (k8\_setfam\_1 \\ & X0 X1)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 (k1\_zfmisc\_1 X1)) \Leftrightarrow (r1\_tarski X0 X1) \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. r1\_tarski X0 X0 \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 X0)) \Rightarrow (k9\_subset\_1 X0 X1 X2 = k3\_xboole\_0 X1 X2) \tag{4}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 X0)) \Rightarrow (m1\_subset\_1 (k9\_subset\_1 X0 X1 X2) (k1\_zfmisc\_1 X0)) \tag{5}$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k1\_zfmisc\_1 X0))) \Rightarrow (m1\_subset\_1 (k8\_setfam\_1 X0 X1) (k1\_zfmisc\_1 X0)) \tag{6}$$

### Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k1\_zfmisc\_1 \\ & X0))) \Rightarrow (\forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k1\_zfmisc\_1 \\ & X0))) \Rightarrow (r1\_tarski (k9\_subset\_1 X0 (k8\_setfam\_1 X0 X1) (k8\_setfam\_1 \\ & X0 X2)) (k8\_setfam\_1 X0 (k9\_subset\_1 (k1\_zfmisc\_1 X0) X1 X2)))) \end{aligned}$$